

NCERT Solutions for Class 10 Maths Unit 8

Introduction to Trigonometry Class 10

Unit 8 Introduction to Trigonometry Exercise 8.1, 8.2, 8.3, 8.4 Solutions

Exercise 8.1 : Solutions of Questions on Page Number : **181**

Q1 :

In ΔABC right angled at B, AB = 24 cm, BC = 7 m. Determine


- (i) $\sin A, \cos A$
- (ii) $\sin C, \cos C$

Answer :

Applying Pythagoras theorem for ΔABC , we obtain


$$\begin{aligned} AC^2 &= AB^2 + BC^2 \\ &= (24 \text{ cm})^2 + (7 \text{ cm})^2 \\ &= (576 + 49) \text{ cm}^2 \\ &= 625 \text{ cm}^2 \end{aligned}$$

$$\therefore AC = \sqrt{625} \text{ cm} = 25 \text{ cm}$$



$$\begin{aligned} \text{(i) } \sin A &= \frac{\text{Side opposite to } \angle A}{\text{Hypotenuse}} = \frac{BC}{AC} \\ &= \frac{7}{25} \end{aligned}$$

$$\begin{aligned} \text{cos A} &= \frac{\text{Side adjacent to } \angle A}{\text{Hypotenuse}} = \frac{AB}{AC} = \frac{24}{25} \\ \text{(ii)} & \end{aligned}$$



$$\sin C = \frac{\text{Side opposite to } \angle C}{\text{Hypotenuse}} = \frac{AB}{AC}$$


$$= \frac{24}{25}$$

$$\cos C = \frac{\text{Side adjacent to } \angle C}{\text{Hypotenuse}} = \frac{BC}{AC}$$

$$= \frac{7}{25}$$

Q2 :

In the given figure find $\tan P - \cot R$



Answer :

Applying Pythagoras theorem for $\triangle PQR$, we obtain


$$PR^2 = PQ^2 + QR^2$$

$$(13 \text{ cm})^2 = (12 \text{ cm})^2 + QR^2$$

$$169 \text{ cm}^2 = 144 \text{ cm}^2 + QR^2$$

$$25 \text{ cm}^2 = QR^2$$

$$= 5 \text{ cm}$$



$$\tan P = \frac{\text{Side opposite to } \angle P}{\text{Side adjacent to } \angle P} = \frac{QR}{PQ}$$

$$= \frac{5}{12}$$

$$\cot R = \frac{\text{Side adjacent to } \angle R}{\text{Side opposite to } \angle R} = \frac{QR}{PQ}$$

$$= \frac{5}{12}$$


$$\tan P - \cot R = \frac{5}{12} - \frac{5}{12} = 0$$

Q3 :

If $\sin A = \frac{3}{4}$, calculate $\cos A$ and $\tan A$.

Answer :

Let ΔABC be a right-angled triangle, right-angled at point B.



Given that,

$$\sin A = \frac{3}{4}$$

$$\frac{BC}{AC} = \frac{3}{4}$$

Let BC be $3k$. Therefore, AC will be $4k$, where k is a positive integer.

Applying Pythagoras theorem in ΔABC , we obtain

$$AC^2 = AB^2 + BC^2$$

$$(4k)^2 = AB^2 + (3k)^2$$

$$16k^2 - 9k^2 = AB^2$$

$$7k^2 = AB^2$$

$$AB = \sqrt{7k}$$

$$\cos A = \frac{\text{Side adjacent to } \angle A}{\text{Hypotenuse}}$$

$$= \frac{AB}{AC} = \frac{\sqrt{7k}}{4k} = \frac{\sqrt{7}}{4}$$

$$\tan A = \frac{\text{Side opposite to } \angle A}{\text{Side adjacent to } \angle A}$$


$$= \frac{BC}{AB} = \frac{3k}{\sqrt{7k}} = \frac{3}{\sqrt{7}}$$

Q4 :

Given $15 \cot A = 8$. Find $\sin A$ and $\sec A$

Answer :

Consider a right-angled triangle, right-angled at B.



$$\cot A = \frac{\text{Side adjacent to } \angle A}{\text{Side opposite to } \angle A}$$

$$= \frac{AB}{BC}$$

It is given that,

$$\cot A = \frac{8}{15}$$

$$\frac{AB}{BC} = \frac{8}{15}$$

Let AB be $8k$. Therefore, BC will be $15k$, where k is a positive integer.

Applying Pythagoras theorem in ΔABC , we obtain

$$AC^2 = AB^2 + BC^2$$

$$= (8k)^2 + (15k)^2$$

$$= 64k^2 + 225k^2$$

$$= 289k^2$$

$$AC = 17k$$

$$\sin A = \frac{\text{Side opposite to } \angle A}{\text{Hypotenuse}} = \frac{BC}{AC}$$

$$= \frac{15k}{17k} = \frac{15}{17}$$

$$\sec A = \frac{\text{Hypotenuse}}{\text{Side adjacent to } \angle A}$$

$$= \frac{AC}{AB} = \frac{17}{8}$$


Q5 :

$$\frac{13}{12}$$

Given $\sec \theta = \frac{13}{12}$, calculate all other trigonometric ratios.

Answer :

Consider a right-angle triangle ΔABC , right-angled at point B.



$$\sec \theta = \frac{\text{Hypotenuse}}{\text{Side adjacent to } \angle \theta}$$

$$\frac{13}{12} = \frac{AC}{AB}$$

If AC is $13k$, AB will be $12k$, where k is a positive integer.

Applying Pythagoras theorem in $\triangle ABC$, we obtain

$$(AC)^2 = (AB)^2 + (BC)^2$$

$$(13k)^2 = (12k)^2 + (BC)^2$$

$$169k^2 = 144k^2 + BC^2$$

$$25k^2 = BC^2$$

$$BC = 5k$$

$$\sin \theta = \frac{\text{Side opposite to } \angle \theta}{\text{Hypotenuse}} = \frac{BC}{AC} = \frac{5k}{13k} = \frac{5}{13}$$

$$\cos \theta = \frac{\text{Side adjacent to } \angle \theta}{\text{Hypotenuse}} = \frac{AB}{AC} = \frac{12k}{13k} = \frac{12}{13}$$

$$\tan \theta = \frac{\text{Side opposite to } \angle \theta}{\text{Side adjacent to } \angle \theta} = \frac{BC}{AB} = \frac{5k}{12k} = \frac{5}{12}$$

$$\cot \theta = \frac{\text{Side adjacent to } \angle \theta}{\text{Side opposite to } \angle \theta} = \frac{AB}{BC} = \frac{12k}{5k} = \frac{12}{5}$$


$$\cosec \theta = \frac{\text{Hypotenuse}}{\text{Side opposite to } \angle \theta} = \frac{AC}{BC} = \frac{13k}{5k} = \frac{13}{5}$$

Q6 :

If $\angle A$ and $\angle B$ are acute angles such that $\cos A = \cos B$, then show that $\angle A = \angle B$.

Answer :

Let us consider a triangle ABC in which $CD \perp AB$.




It is given that \cos

$$A = \cos B$$

$$\Rightarrow \frac{AD}{AC} = \frac{BD}{BC} \dots (1)$$

We have to prove $\angle A = \angle B$. To prove this, let us extend AC to P such that BC = CP.



From equation (1), we obtain

$$\begin{aligned} \frac{AD}{BD} &= \frac{AC}{BC} \\ \Rightarrow \frac{AD}{BD} &= \frac{AC}{CP} \quad (\text{By construction, we have } BC = CP) \end{aligned} \dots (2)$$

By using the converse of B.P.T,

$$CD \parallel BP$$

$\Rightarrow \angle ACD = \angle CPB$ (Corresponding angles) ... (3) And,

$\angle BCD = \angle CBP$ (Alternate interior angles) ... (4) By

construction, we have $BC = CP$.

$\therefore \angle CPB = \angle CBP$ (Angle opposite to equal sides of a triangle) ... (5)

From equations (3), (4), and (5), we obtain

$$\angle ACD = \angle BCD \dots (6)$$

In $\triangle CAD$ and $\triangle CBD$,

$\angle ACD = \angle BCD$ [Using equation (6)]

$\angle CDA = \angle CDB$ [Both 90°]


Therefore, the remaining angles should be equal.

$$\therefore \angle CAD = \angle CBD$$

$$\Rightarrow \angle A = \angle B$$

Alternatively,

Let us consider a triangle ABC in which $CD \perp AB$.



It is given that,

$$\cos A = \cos B$$

$$\Rightarrow \frac{AD}{AC} = \frac{BD}{BC}$$

$$\Rightarrow \frac{AD}{BD} = \frac{AC}{BC}$$

$$\text{Let } \frac{AD}{BD} = \frac{AC}{BC} = k$$

$$\Rightarrow AD = k BD \dots (1)$$

$$\text{And, } AC = k BC \dots (2)$$

Using Pythagoras theorem for triangles CAD and CBD, we obtain

$$CD^2 = AC^2 - AD^2 \dots (3)$$

$$\text{And, } CD^2 = BC^2 - BD^2 \dots (4)$$

From equations (3) and (4), we obtain

$$AC^2 - AD^2 = BC^2 - BD^2 \Rightarrow (k$$

$$BC)^2 - (k BD)^2 = BC^2 - BD^2 \Rightarrow k^2 =$$

$$(BC^2 - BD^2) = BC^2 - BD^2 \Rightarrow k^2 =$$

$$1 \Rightarrow k = 1$$

Putting this value in equation (2), we obtain

$$AC = BC$$

$\Rightarrow \angle A = \angle B$ (Angles opposite to equal sides of a triangle)


Q7 :

If $\cot \theta = \frac{7}{8}$, evaluate

$$(i) \frac{(1+\sin\theta)(1-\sin\theta)}{(1+\cos\theta)(1-\cos\theta)} \quad (ii) \cot^2 \theta$$

Answer :

Let us consider a right triangle ABC, right-angled at point B.



$$\cot \theta = \frac{\text{Side adjacent to } \angle \theta}{\text{Side opposite to } \angle \theta} = \frac{BC}{AB}$$

$$= \frac{7}{8}$$

If BC is $7k$, then AB will be $8k$, where k is a positive integer.

Applying Pythagoras theorem in ΔABC , we obtain

$$\begin{aligned} AC^2 &= AB^2 + BC^2 \\ &= (8k)^2 + (7k)^2 \\ &= 64k^2 + 49k^2 \end{aligned}$$

$$= 113k^2$$

$$AC = \sqrt{113}k$$

$$\sin \theta = \frac{\text{Side opposite to } \angle \theta}{\text{Hypotenuse}} = \frac{AB}{AC}$$

$$= \frac{8k}{\sqrt{113}k} = \frac{8}{\sqrt{113}}$$

$$\cos \theta = \frac{\text{Side adjacent to } \angle \theta}{\text{Hypotenuse}} = \frac{BC}{AC}$$

$$= \frac{7k}{\sqrt{113}k} = \frac{7}{\sqrt{113}}$$

$$(i) \frac{(1+\sin \theta)(1-\sin \theta)}{(1+\cos \theta)(1-\cos \theta)} = \frac{(1-\sin^2 \theta)}{(1-\cos^2 \theta)}$$

$$= \frac{1 - \left(\frac{8}{\sqrt{113}}\right)^2}{1 - \left(\frac{7}{\sqrt{113}}\right)^2} = \frac{1 - \frac{64}{113}}{1 - \frac{49}{113}}$$

$$= \frac{\frac{49}{113}}{\frac{64}{113}} = \frac{49}{64}$$

$$(ii) \cot^2 \theta = (\cot \theta)^2 = \left(\frac{7}{8}\right)^2 = \frac{49}{64}$$

Q8 :

$$\frac{1 - \tan^2 A}{1 + \tan^2 A} = \cos^2 A - \sin^2 A \text{ or not.}$$


If $3 \cot A = 4$, Check whether

Answer :

It is given that $3\cot A = 4$

$$\text{Or, } \cot A = \frac{4}{3}$$

Consider a right triangle ABC, right-angled at point B.



$$\cot A = \frac{\text{Side adjacent to } \angle A}{\text{Side opposite to } \angle A}$$

$$\frac{AB}{BC} = \frac{4}{3}$$

If AB is $4k$, then BC will be $3k$, where k is a positive integer.

In $\triangle ABC$,

$$(AC)^2 = (AB)^2 + (BC)^2$$

$$\begin{aligned} &= (4k)^2 + (3k)^2 \\ &= 16k^2 + 9k^2 \end{aligned}$$

$$= 25k^2$$

$$AC = 5k$$

$$\cos A = \frac{\text{Side adjacent to } \angle A}{\text{Hypotenuse}} = \frac{AB}{AC}$$

$$= \frac{4k}{5k} = \frac{4}{5}$$

$$\sin A = \frac{\text{Side opposite to } \angle A}{\text{Hypotenuse}} = \frac{BC}{AC}$$

$$= \frac{3k}{5k} = \frac{3}{5}$$

$$\tan A = \frac{\text{Side opposite to } \angle A}{\text{Hypotenuse}} = \frac{BC}{AB}$$

$$= \frac{3k}{4k} = \frac{3}{4}$$

$$\frac{1 - \tan^2 A}{1 + \tan^2 A} = \frac{1 - \left(\frac{3}{4}\right)^2}{1 + \left(\frac{3}{4}\right)^2} = \frac{1 - \frac{9}{16}}{1 + \frac{9}{16}}$$

$$= \frac{\frac{7}{16}}{\frac{25}{16}} = \frac{7}{25}$$

$$\cos^2 A - \sin^2 A = \left(\frac{4}{5}\right)^2 - \left(\frac{3}{5}\right)^2$$

$$= \frac{16}{25} - \frac{9}{25} = \frac{7}{25}$$

$$\therefore \frac{1 - \tan^2 A}{1 + \tan^2 A} = \cos^2 A - \sin^2 A$$


Q9 :

$$\tan A = \frac{1}{\sqrt{3}}$$

In ΔABC , right angled at B. If

- (i) $\sin A \cos C + \cos A \sin C$
- (ii) $\cos A \cos C - \sin A \sin C$

Answer :



$$\tan A = \frac{1}{\sqrt{3}}$$

$$\frac{BC}{AB} = \frac{1}{\sqrt{3}}$$

If BC is k , then AB will be $\sqrt{3}k$, where k is a positive integer.

In $\triangle ABC$,

$$AC^2 = AB^2 + BC^2$$

$$= (\sqrt{3}k)^2 + (k)^2$$

$$= 3k^2 + k^2 = 4k^2$$

$$AC = 2k$$

$$\sin A = \frac{\text{Side opposite to } \angle A}{\text{Hypotenuse}} = \frac{BC}{AC} = \frac{k}{2k} = \frac{1}{2}$$

$$\cos A = \frac{\text{Side adjacent to } \angle A}{\text{Hypotenuse}} = \frac{AB}{AC} = \frac{\sqrt{3}k}{2k} = \frac{\sqrt{3}}{2}$$

$$\sin C = \frac{\text{Side opposite to } \angle C}{\text{Hypotenuse}} = \frac{AB}{AC} = \frac{\sqrt{3}k}{2k} = \frac{\sqrt{3}}{2}$$

$$\cos C = \frac{\text{Side adjacent to } \angle C}{\text{Hypotenuse}} = \frac{BC}{AC} = \frac{k}{2k} = \frac{1}{2}$$

(i) $\sin A \cos C + \cos A \sin C$

$$= \left(\frac{1}{2}\right)\left(\frac{1}{2}\right) + \left(\frac{\sqrt{3}}{2}\right)\left(\frac{\sqrt{3}}{2}\right) = \frac{1}{4} + \frac{3}{4}$$

$$= \frac{4}{4} = 1$$

(ii) $\cos A \cos C - \sin A \sin C$

$$= \left(\frac{\sqrt{3}}{2}\right)\left(\frac{1}{2}\right) - \left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right) = \frac{\sqrt{3}}{4} - \frac{\sqrt{3}}{4} = 0$$

Q10 :

In ΔPQR , right angled at Q, $PR + QR = 25$ cm and $PQ = 5$ cm. Determine the values of $\sin P$, $\cos P$ and $\tan P$.


Answer :

Given that, $PR + QR = 25$

$$PQ = 5$$

Let PR be x .

Therefore, $QR = 25 - x$



Applying Pythagoras theorem in ΔPQR , we obtain

$$PR^2 = PQ^2 + QR^2$$

$$x^2 = (5)^2 + (25 - x)^2 \quad x^2 =$$

$$25 + 625 + x^2 - 50x$$

$$50x = 650$$

$$x = 13$$

Therefore, $PR = 13$ cm

$$QR = (25 - 13) \text{ cm} = 12 \text{ cm}$$

$$\sin P = \frac{\text{Side opposite to } \angle P}{\text{Hypotenuse}} = \frac{QR}{PR} = \frac{12}{13}$$

$$\cos P = \frac{\text{Side adjacent to } \angle P}{\text{Hypotenuse}} = \frac{PQ}{PR} = \frac{5}{13}$$

$$\tan P = \frac{\text{Side opposite to } \angle P}{\text{Side adjacent to } \angle P} = \frac{QR}{PQ} = \frac{12}{5}$$

Q11 :

State whether the following are true or false. Justify your answer.

(i) The value of $\tan A$ is always less than 1.

$$\frac{12}{5}$$

(ii) $\sec A = \frac{4}{3}$ for some value of angle A.

(iii) $\cos A$ is the abbreviation used for the cosecant of angle A.


(iv) $\cot A$ is the product of cot and A

$$\frac{4}{3}$$

(v) $\sin \theta = \frac{4}{3}$, for some angle θ

Answer :

(i) Consider a $\triangle ABC$, right-angled at B.



$$\tan A = \frac{\text{Side opposite to } \angle A}{\text{Side adjacent to } \angle A}$$

$$= \frac{12}{5}$$


$$\text{But } \frac{12}{5} > 1$$

$$\therefore \tan A > 1$$

So, $\tan A < 1$ is not always true.

Hence, the given statement is false.

$$(ii) \sec A = \frac{12}{5}$$



$$\frac{\text{Hypotenuse}}{\text{Side adjacent to } \angle A} = \frac{12}{5}$$

$$\frac{AC}{AB} = \frac{12}{5}$$

Let AC be $12k$, AB will be $5k$, where k is a positive integer.

Applying Pythagoras theorem in ΔABC , we obtain

$$AC^2 = AB^2 + BC^2$$

$$(12k)^2 = (5k)^2 + BC^2$$

$$144k^2 = 25k^2 + BC^2$$

$$BC^2 = 119k^2$$

$$BC = 10.9k$$

It can be observed that for given two sides $AC = 12k$ and $AB = 5k$,

BC should be such that,

$$AC - AB < BC < AC + AB$$

$$12k - 5k < BC < 12k + 5k$$

$$7k < BC < 17k$$

However, $BC = 10.9k$. Clearly, such a triangle is possible and hence, such value of $\sec A$ is possible.

Hence, the given statement is true.

(iii) Abbreviation used for cosecant of angle A is cosec A. And cos A is the abbreviation used for cosine of angle A.

Hence, the given statement is false.

(iv) $\cot A$ is not the product of \cot and A. It is the cotangent of $\angle A$.

Hence, the given statement is false.

$$(v) \sin \theta = \frac{4}{3}$$

We know that in a right-angled triangle,

$$\sin \theta = \frac{\text{Side opposite to } \angle \theta}{\text{Hypotenuse}}$$

In a right-angled triangle, hypotenuse is always greater than the remaining two sides. Therefore, such value of $\sin \theta$ is not possible.

Hence, the given statement is false

Exercise 8.2 : Solutions of Questions on Page Number : 187

Q1 :

Evaluate the following

$$(i) \sin 60^\circ \cos 30^\circ + \sin 30^\circ \cos 60^\circ$$

$$(ii) 2\tan^2 45^\circ + \cos^2 30^\circ - \sin^2 60^\circ$$

$$(iii) \frac{\cos 45^\circ}{\sec 30^\circ + \operatorname{cosec} 30^\circ}$$

$$(iv) \frac{\sin 30^\circ + \tan 45^\circ - \operatorname{cosec} 60^\circ}{\sec 30^\circ + \cos 60^\circ + \cot 45^\circ}$$

$$(v) \frac{5\cos^2 60^\circ + 4\sec^2 30^\circ - \tan^2 45^\circ}{\sin^2 30^\circ + \cos^2 30^\circ}$$

Answer :

$$(i) \sin 60^\circ \cos 30^\circ + \sin 30^\circ \cos 60^\circ$$

$$\begin{aligned} &= \left(\frac{\sqrt{3}}{2}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(\frac{1}{2}\right)\left(\frac{1}{2}\right) \\ &= \frac{3}{4} + \frac{1}{4} = \frac{4}{4} = 1 \end{aligned}$$

$$(ii) 2\tan^2 45^\circ + \cos^2 30^\circ - \sin^2 60^\circ$$

$$\begin{aligned} &= 2(1)^2 + \left(\frac{\sqrt{3}}{2}\right)^2 - \left(\frac{\sqrt{3}}{2}\right)^2 \\ &= 2 + \frac{3}{4} - \frac{3}{4} = 2 \end{aligned}$$

$$(iii) \frac{\cos 45^\circ}{\sec 30^\circ + \operatorname{cosec} 30^\circ}$$

$$\begin{aligned}
 &= \frac{\frac{1}{\sqrt{2}}}{\frac{2}{\sqrt{3}} + 2} = \frac{\frac{1}{\sqrt{2}}}{\frac{2+2\sqrt{3}}{\sqrt{3}}} \\
 &= \frac{\sqrt{3}}{\sqrt{2}(2+2\sqrt{3})} = \frac{\sqrt{3}}{2\sqrt{2}+2\sqrt{6}} \\
 &= \frac{\sqrt{3}(2\sqrt{6}-2\sqrt{2})}{(2\sqrt{6}+2\sqrt{2})(2\sqrt{6}-2\sqrt{2})} \\
 &= \frac{2\sqrt{3}(\sqrt{6}-\sqrt{2})}{(2\sqrt{6})^2 - (2\sqrt{2})^2} = \frac{2\sqrt{3}(\sqrt{6}-\sqrt{2})}{24-8} = \frac{2\sqrt{3}(\sqrt{6}-\sqrt{2})}{16} \\
 &= \frac{\sqrt{18}-\sqrt{6}}{8} = \frac{3\sqrt{2}-\sqrt{6}}{8}
 \end{aligned}$$

$$\frac{\sin 30^\circ + \tan 45^\circ - \operatorname{cosec} 60^\circ}{\sec 30^\circ + \cos 60^\circ + \cot 45^\circ}$$

$$\begin{aligned}
 &= \frac{\frac{1}{2} + 1 - \frac{2}{\sqrt{3}}}{\frac{2}{\sqrt{3}} + \frac{1}{2} + 1} = \frac{\frac{3}{2} - \frac{2}{\sqrt{3}}}{\frac{3}{2} + \frac{2}{\sqrt{3}}}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\frac{3\sqrt{3}-4}{2\sqrt{3}}}{\frac{3\sqrt{3}+4}{2\sqrt{3}}} = \frac{(3\sqrt{3}-4)}{(3\sqrt{3}+4)}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{(3\sqrt{3}-4)(3\sqrt{3}-4)}{(3\sqrt{3}+4)(3\sqrt{3}-4)} = \frac{(3\sqrt{3}-4)^2}{(3\sqrt{3})^2 - (4)^2}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{27+16-24\sqrt{3}}{27-16} = \frac{43-24\sqrt{3}}{11}
 \end{aligned}$$

$$\begin{aligned}
 &(v) \quad \frac{5\cos^2 60^\circ + 4\sec^2 30^\circ - \tan^2 45^\circ}{\sin^2 30^\circ + \cos^2 30^\circ}
 \end{aligned}$$

$$= \frac{5\left(\frac{1}{2}\right)^2 + 4\left(\frac{2}{\sqrt{3}}\right)^2 - (1)^2}{\left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2}$$

$$= \frac{5\left(\frac{1}{4}\right) + \left(\frac{16}{3}\right) - 1}{\frac{1}{4} + \frac{3}{4}}$$

$$= \frac{15 + 64 - 12}{12} = \frac{67}{12}$$

Q2 :

Choose the correct option and justify your choice.

(i) $\frac{2 \tan 30^\circ}{1 + \tan^2 30^\circ} =$

(A). $\sin 60^\circ$

(B). $\cos 60^\circ$

(C). $\tan 60^\circ$

(D). $\sin 30^\circ$

(ii) $\frac{1 - \tan^2 45^\circ}{1 + \tan^2 45^\circ} =$

(A). $\tan 90^\circ$ (B). $\frac{1}{1}$

(C). $\sin 45^\circ$

(D). 0

(iii) $\sin 2A = 2 \sin A$ is true when $A =$

(A). 0°

(B). 30°

(C). 45°

(D). 60°

(iv) $\frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ} =$

(A). $\cos 60^\circ$

(B). $\sin 60^\circ$ (C).
 $\tan 60^\circ$

(D). $\sin 30^\circ$

Answer :

$$(i) \frac{2 \tan 30^\circ}{1 + \tan^2 30^\circ}$$

$$= \frac{2 \left(\frac{1}{\sqrt{3}} \right)}{1 + \left(\frac{1}{\sqrt{3}} \right)^2} = \frac{\frac{2}{\sqrt{3}}}{1 + \frac{1}{3}} = \frac{\frac{2}{\sqrt{3}}}{\frac{4}{3}} \\ = \frac{6}{4\sqrt{3}} = \frac{\sqrt{3}}{2}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

Out of the given alternatives, only

Hence, (A) is correct.

$$(ii) \frac{1 - \tan^2 45^\circ}{1 + \tan^2 45^\circ}$$

$$= \frac{1 - (1)^2}{1 + (1)^2} = \frac{1 - 1}{1 + 1} = \frac{0}{2} = 0$$

Hence, (D) is correct.

(iii) Out of the given alternatives, only $A = 0^\circ$ is correct.

As $\sin 2A = \sin 0^\circ = 0$

$$2 \sin A = 2 \sin 0^\circ = 2(0) = 0$$

Hence, (A) is correct.

$$(iv) \frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ}$$

$$= \frac{2 \left(\frac{1}{\sqrt{3}} \right)}{1 - \left(\frac{1}{\sqrt{3}} \right)^2} = \frac{\frac{2}{\sqrt{3}}}{1 - \frac{1}{3}} = \frac{\frac{2}{\sqrt{3}}}{\frac{2}{3}} \\ = \sqrt{3}$$

Out of the given alternatives, only $\tan 60^\circ = \sqrt{3}$ Hence,
(C) is correct.

Q3 :

$$\text{If } \tan(A + B) = \sqrt{3} \text{ and } \tan(A - B) = \frac{1}{\sqrt{3}};$$

$0^\circ < A + B \leq 90^\circ$, $A > B$ find A and B.

Answer :

$$\tan(A + B) = \sqrt{3}$$

$$\Rightarrow \tan(A + B) = \tan 60$$

$$\Rightarrow A + B = 60 \dots (1)$$

$$\tan(A - B) = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \tan(A - B) = \tan 30$$

$$\Rightarrow A - B = 30 \dots (2)$$

On adding both equations, we obtain

$$2A = 90$$

$$\Rightarrow A = 45$$

From equation (1), we obtain

$$45 + B = 60$$

$$B = 15$$

Therefore, $\angle A = 45^\circ$ and $\angle B = 15^\circ$

Q4 :

State whether the following are true or false. Justify your answer.

- (i) $\sin(A + B) = \sin A + \sin B$
- (ii) The value of $\sin A$ increases as A increases
- (iii) The value of $\cos A$ increases as A increases
- (iv) $\sin A = \cos A$ for all values of A ,
- (v) $\cot A$ is not defined for $A = 0^\circ$

Answer :

(i) $\sin(A + B) = \sin A + \sin B$ Let $A = 30^\circ$ and $B = 60^\circ$ $\sin(A + B) = \sin(30^\circ + 60^\circ)$

$$= \sin 90^\circ =$$

1

$$\sin A + \sin B = \sin 30^\circ + \sin 60^\circ$$

$$= \frac{1}{2} + \frac{\sqrt{3}}{2} = \frac{1+\sqrt{3}}{2}$$

Clearly, $\sin(A + B) \neq \sin A + \sin B$

Hence, the given statement is false.

(ii) The value of $\sin \theta$ increases as θ increases in the interval of $0^\circ < \theta < 90^\circ$ as $\sin 0^\circ = 0$

$$\sin 30^\circ = \frac{1}{2} = 0.5$$

$$\sin 45^\circ = \frac{1}{\sqrt{2}} = 0.707$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2} = 0.866$$

$$\sin 90^\circ = 1$$

Hence, the given statement is true.

(iii) $\cos 0^\circ = 1$

$$\cos 30^\circ = \frac{\sqrt{3}}{2} = 0.866$$

$$\cos 45^\circ = \frac{1}{\sqrt{2}} = 0.707$$

$$\cos 60^\circ = \frac{1}{2} = 0.5$$

$$\cos 90^\circ = 0$$

It can be observed that the value of $\cos \theta$ does not increase in the interval of $0^\circ < \theta < 90^\circ$.

Hence, the given statement is false.

(iv) $\sin \theta = \cos \theta$ for all values of θ .

This is true when $\theta = 45^\circ$

As $\sin 45^\circ = \frac{1}{\sqrt{2}}$

$$\cos 45^\circ = \frac{1}{\sqrt{2}}$$

It is not true for all other values of θ .

As $\sin 30^\circ = \frac{1}{2}$ and $\cos 30^\circ = \frac{\sqrt{3}}{2}$,

Hence, the given statement is false.

(v) $\cot A$ is not defined for $A = 0^\circ$

As $\cot A = \frac{\cos A}{\sin A}$,

$$\cot 0^\circ = \frac{\cos 0^\circ}{\sin 0^\circ} = \frac{1}{0} = \text{undefined}$$

Hence, the given statement is true.

Exercise 8.3 : Solutions of Questions on Page Number : 189

Q1 :

Evaluate

$$\frac{\sin 18^\circ}{\cos 72^\circ} \quad (\text{I})$$

$$\frac{\tan 26^\circ}{\cot 64^\circ} \quad (\text{II})$$

(III) $\cos 48^\circ - \sin 42^\circ$

(IV) $\cosec 31^\circ - \sec 59^\circ$

Answer :

$$(\text{I}) \frac{\sin 18^\circ}{\cos 72^\circ} = \frac{\sin(90^\circ - 72^\circ)}{\cos 72^\circ}$$

$$= \frac{\cos 72^\circ}{\cos 72^\circ} = 1$$

$$(\text{II}) \frac{\tan 26^\circ}{\cot 64^\circ} = \frac{\tan(90^\circ - 64^\circ)}{\cot 64^\circ}$$

$$= \frac{\cot 64^\circ}{\cot 64^\circ} = 1$$

$$(III) \cos 48^\circ - \sin 42^\circ = \cos (90^\circ - 42^\circ) - \sin 42^\circ$$

$$= \sin 42^\circ - \sin 42^\circ$$

$$= 0$$

$$(IV) \operatorname{cosec} 31^\circ - \sec 59^\circ = \operatorname{cosec} (90^\circ - 59^\circ) - \sec 59^\circ$$

$$= \sec 59^\circ - \sec 59^\circ$$

$$= 0$$

Q2 :

Show that

$$(I) \tan 48^\circ \tan 23^\circ \tan 42^\circ \tan 67^\circ = 1$$

$$(II) \cos 38^\circ \cos 52^\circ - \sin 38^\circ \sin 52^\circ = 0$$

Answer :

$$(I) \tan 48^\circ \tan 23^\circ \tan 42^\circ \tan 67^\circ$$

$$= \tan (90^\circ - 42^\circ) \tan (90^\circ - 67^\circ) \tan 42^\circ \tan 67^\circ$$

$$= \cot 42^\circ \cot 67^\circ \tan 42^\circ \tan 67^\circ$$

$$= (\cot 42^\circ \tan 42^\circ) (\cot 67^\circ \tan 67^\circ)$$

$$= (1) (1)$$

$$= 1$$

$$(II) \cos 38^\circ \cos 52^\circ - \sin 38^\circ \sin 52^\circ$$

$$= \cos (90^\circ - 52^\circ) \cos (90^\circ - 38^\circ) - \sin 38^\circ \sin 52^\circ$$

$$= \sin 52^\circ \sin 38^\circ - \sin 38^\circ \sin 52^\circ$$

$$= 0$$

Q3 :

If $\tan 2A = \cot (A - 18^\circ)$, where $2A$ is an acute angle, find the value of A .

Answer : Given that, \tan

$$2A = \cot (A - 18^\circ) \cot (90^\circ -$$

$$2A) = \cot (A - 18^\circ) 90^\circ - 2A$$

$$= A - 18^\circ$$

$$108^\circ = 3A$$

$$A = 36^\circ$$

Q4 :

If $\tan A = \cot B$, prove that $A + B = 90^\circ$

Answer : Given

that, $\tan A = \cot B$

$$\tan A = \tan (90^\circ - B)$$

$$A = 90^\circ - B$$

$$A + B = 90^\circ$$

Q5 :

If $\sec 4A = \operatorname{cosec} (A - 20^\circ)$, where $4A$ is an acute angle, find the value of A .

Answer : Given that, $\sec 4A =$

$$\operatorname{cosec} (A - 20^\circ) \operatorname{cosec} (90^\circ - 4A) =$$

$$\operatorname{cosec} (A - 20^\circ)$$

$$90^\circ - 4A = A - 20^\circ$$

$$110^\circ = 5A$$

$$A = 22^\circ$$

Q6 :

If A , B and C are interior angles of a triangle ABC then show that

$$\sin\left(\frac{B+C}{2}\right) = \cos\frac{A}{2}$$

Answer :

We know that for a triangle ABC ,

$$\angle A + \angle B + \angle C = 180^\circ$$

$$\square \quad \angle B + \angle C = 180^\circ - \angle A$$

$$\begin{aligned}\frac{\angle B + \angle C}{2} &= 90^\circ - \frac{\angle A}{2} \\ \sin\left(\frac{\angle B + \angle C}{2}\right) &= \sin\left(90^\circ - \frac{\angle A}{2}\right) \\ &= \cos\left(\frac{\angle A}{2}\right)\end{aligned}$$

Q7 :

Express $\sin 67^\circ + \cos 75^\circ$ in terms of trigonometric ratios of angles between 0° and 45° .

Answer :

$$\begin{aligned}\sin 67^\circ + \cos 75^\circ &\\ &= \sin (90^\circ - 23^\circ) + \cos (90^\circ - 15^\circ) \\ &= \cos 23^\circ + \sin 15^\circ\end{aligned}$$

Exercise 8.4 : Solutions of Questions on Page Number : 193

Q1 :

Express the trigonometric ratios $\sin A$, $\sec A$ and $\tan A$ in terms of $\cot A$.

Answer :

We know that,

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\frac{1}{\operatorname{cosec}^2 A} = \frac{1}{1 + \cot^2 A}$$

$$\sin^2 A = \frac{1}{1 + \cot^2 A}$$

$$\sin A = \pm \frac{1}{\sqrt{1 + \cot^2 A}}$$

$\sqrt{1 + \cot^2 A}$ will always be positive as we are adding two positive quantities.

$$\sin A = \frac{1}{\sqrt{1 + \cot^2 A}}$$

Therefore,

$$\tan A = \frac{\sin A}{\cos A}$$

We know that,

www.eduinput.com

$$\cot A = \frac{\cos A}{\sin A}$$

However,

$$\tan A = \frac{1}{\cot A}$$

Therefore,

$$\text{Also, } \sec^2 A = 1 + \tan^2 A$$

$$= 1 + \frac{1}{\cot^2 A}$$

$$= \frac{\cot^2 A + 1}{\cot^2 A}$$

$$\sec A = \frac{\sqrt{\cot^2 A + 1}}{\cot A}$$

Q2 :

Write all the other trigonometric ratios of $\angle A$ in terms of $\sec A$.

Answer :

We know that,

$$\cos A = \frac{1}{\sec A}$$

$$\text{Also, } \sin^2 A + \cos^2 A = 1 \quad \sin^2$$

$$A = 1 - \cos^2 A$$

$$\begin{aligned} \sin A &= \sqrt{1 - \left(\frac{1}{\sec A}\right)^2} \\ &= \sqrt{\frac{\sec^2 A - 1}{\sec^2 A}} = \frac{\sqrt{\sec^2 A - 1}}{\sec A} \end{aligned}$$

$$\tan^2 A + 1 = \sec^2 A \tan^2 A$$

$$= \sec^2 A - 1$$

$$\tan A = \sqrt{\sec^2 A - 1}$$

$$\cot A = \frac{\cos A}{\sin A} = \frac{1}{\frac{\sec A}{\sqrt{\sec^2 A - 1}}} \\ = \frac{1}{\sec A}$$

$$\operatorname{cosec} A = \frac{1}{\sin A} = \frac{\sec A}{\sqrt{\sec^2 A - 1}}$$

Q3 :

Evaluate

$$(i) \frac{\sin^2 63^\circ + \sin^2 27^\circ}{\cos^2 17^\circ + \cos^2 73^\circ}$$

$$(ii) \sin 25^\circ \cos 65^\circ + \cos 25^\circ \sin 65^\circ$$

Answer :

$$(i) \frac{\sin^2 63^\circ + \sin^2 27^\circ}{\cos^2 17^\circ + \cos^2 73^\circ} \\ = \frac{[\sin(90^\circ - 27^\circ)]^2 + \sin^2 27^\circ}{[\cos(90^\circ - 73^\circ)]^2 + \cos^2 73^\circ} \\ = \frac{[\cos 27^\circ]^2 + \sin^2 27^\circ}{[\sin 73^\circ]^2 + \cos^2 73^\circ} \\ = \frac{\cos^2 27^\circ + \sin^2 27^\circ}{\sin^2 73^\circ + \cos^2 73^\circ} \\ = \frac{1}{1} \quad (\text{As } \sin^2 A + \cos^2 A = 1)$$

$$= 1$$

$$(ii) \sin 25^\circ \cos 65^\circ + \cos 25^\circ \sin 65^\circ$$

$$= (\sin 25^\circ) \{ \cos(90^\circ - 25^\circ) \} + \cos 25^\circ \{ \sin(90^\circ - 25^\circ) \} \\ = (\sin 25^\circ)(\sin 25^\circ) + (\cos 25^\circ)(\cos 25^\circ)$$

$$= \sin^2 25^\circ + \cos^2 25^\circ \\ = 1 \quad (\text{As } \sin^2 A + \cos^2 A = 1)$$

Q4 :

Choose the correct option. Justify your choice.

(i) $9 \sec^2 A - 9 \tan^2 A =$

- (A) 1
- (B) 9
- (C) 8
- (D) 0

(ii) $(1 + \tan \theta + \sec \theta)(1 + \cot \theta - \operatorname{cosec} \theta)$

- (A) 0
- (B) 1
- (C) 2
- (D) -1

(iii) $(\sec A + \tan A)(1 - \sin A) =$

- (A) $\sec A$
- (B) $\sin A$
- (C) $\operatorname{cosec} A$
- (D) $\cos A$

(iv)
$$\frac{1 + \tan^2 A}{1 + \cot^2 A}$$

- (A) $\sec^2 A$
- (B) -1
- (C) $\cot^2 A$
- (D) $\tan^2 A$

Answer :

(i) $9 \sec^2 A - 9 \tan^2 A$

$= 9 (\sec^2 A - \tan^2 A)$

$= 9 (1) [\text{As } \sec^2 A - \tan^2 A = 1]$

$= 9$

Hence, alternative (B) is correct.

(ii)
 $(1 + \tan \theta + \sec \theta)(1 + \cot \theta - \operatorname{cosec} \theta)$

www.eduinput.com

$$\begin{aligned}
 &= \left(1 + \frac{\sin \theta}{\cos \theta} + \frac{1}{\cos \theta}\right) \left(1 + \frac{\cos \theta}{\sin \theta} - \frac{1}{\sin \theta}\right) \\
 &= \left(\frac{\cos \theta + \sin \theta + 1}{\cos \theta}\right) \left(\frac{\sin \theta + \cos \theta - 1}{\sin \theta}\right) \\
 &= \frac{(\sin \theta + \cos \theta)^2 - (1)^2}{\sin \theta \cos \theta} \\
 &= \frac{\sin^2 \theta + \cos^2 \theta + 2 \sin \theta \cos \theta - 1}{\sin \theta \cos \theta} \\
 &= \frac{1 + 2 \sin \theta \cos \theta - 1}{\sin \theta \cos \theta} \\
 &= \frac{2 \sin \theta \cos \theta}{\sin \theta \cos \theta} = 2
 \end{aligned}$$

Hence, alternative (C) is correct.

(iii) $(\sec A + \tan A)(1 - \sin A)$

$$\begin{aligned}
 &= \left(\frac{1}{\cos A} + \frac{\sin A}{\cos A}\right)(1 - \sin A) \\
 &= \left(\frac{1 + \sin A}{\cos A}\right)(1 - \sin A) \\
 &= \frac{1 - \sin^2 A}{\cos A} = \frac{\cos^2 A}{\cos A}
 \end{aligned}$$

$= \cos A$

Hence, alternative (D) is correct.

$$\frac{1 + \tan^2 A}{1 + \cot^2 A} = \frac{1 + \frac{\sin^2 A}{\cos^2 A}}{1 + \frac{\cos^2 A}{\sin^2 A}}$$

$$\begin{aligned}
 &= \frac{\frac{\cos^2 A + \sin^2 A}{\cos^2 A}}{\frac{\sin^2 A + \cos^2 A}{\sin^2 A}} = \frac{\frac{1}{\cos^2 A}}{\frac{1}{\sin^2 A}} \\
 &= \frac{\sin^2 A}{\cos^2 A} = \tan^2 A
 \end{aligned}$$

Hence, alternative (D) is correct.

Q5 :

Prove the following identities, where the angles involved are acute angles for which the expressions are defined.

Answer :

$$(i) (\csc \theta - \cot \theta)^2 = \frac{1 - \cos \theta}{1 + \cos \theta}$$

$$\begin{aligned} \text{L.H.S.} &= (\csc \theta - \cot \theta)^2 \\ &= \left(\frac{1}{\sin \theta} - \frac{\cos \theta}{\sin \theta} \right)^2 \\ &= \frac{(1 - \cos \theta)^2}{(\sin \theta)^2} = \frac{(1 - \cos \theta)^2}{\sin^2 \theta} \\ &= \frac{(1 - \cos \theta)^2}{1 - \cos^2 \theta} = \frac{(1 - \cos \theta)^2}{(1 - \cos \theta)(1 + \cos \theta)} = \frac{1 - \cos \theta}{1 + \cos \theta} \\ &= \text{R.H.S.} \end{aligned}$$

$$(ii) \frac{\cos A}{1 + \sin A} + \frac{1 + \sin A}{\cos A} = 2 \sec A$$

$$\begin{aligned} \text{L.H.S.} &= \frac{\cos A}{1 + \sin A} + \frac{1 + \sin A}{\cos A} \\ &= \frac{\cos^2 A + (1 + \sin A)^2}{(1 + \sin A)(\cos A)} \\ &= \frac{\cos^2 A + 1 + \sin^2 A + 2\sin A}{(1 + \sin A)(\cos A)} \\ &= \frac{\sin^2 A + \cos^2 A + 1 + 2\sin A}{(1 + \sin A)(\cos A)} \\ &= \frac{1 + 1 + 2\sin A}{(1 + \sin A)(\cos A)} = \frac{2 + 2\sin A}{(1 + \sin A)(\cos A)} \\ &= \frac{2(1 + \sin A)}{(1 + \sin A)(\cos A)} = \frac{2}{\cos A} = 2 \sec A \\ &= \text{R.H.S.} \end{aligned}$$

$$(iii) \frac{\tan \theta}{1 - \cot \theta} + \frac{\cot \theta}{1 - \tan \theta} = 1 + \sec \theta \csc \theta$$

www.eduinput.com

$$\begin{aligned} \text{L.H.S.} &= \frac{\tan \theta}{1 - \cot \theta} + \frac{\cot \theta}{1 - \tan \theta} \\ &= \frac{\frac{\sin \theta}{\cos \theta}}{1 - \frac{\cos \theta}{\sin \theta}} + \frac{\frac{\cos \theta}{\sin \theta}}{1 - \frac{\sin \theta}{\cos \theta}} \\ &= \frac{\sin \theta}{\sin \theta - \cos \theta} + \frac{\cos \theta}{\cos \theta - \sin \theta} \\ &= \frac{\sin \theta}{\cos \theta(\sin \theta - \cos \theta)} - \frac{\cos \theta}{\sin \theta(\sin \theta - \cos \theta)} \end{aligned}$$