



www.eduinput.com

NCERT Solutions for Class 11 Maths Chapter 12

Introduction to Three Dimensional Geometry Class 11

Chapter 12 Introduction to Three Dimensional Geometry Exercise 12.1, 12.2, 12.3, miscellaneous Solutions

Exercise 12.1 : Solutions of Questions on Page Number : 271

Q1 :

A point is on the x -axis. What are its y -coordinates and z -coordinates?

Answer :

If a point is on the x -axis, then its y -coordinates and z -coordinates are zero.

Q2 :

A point is in the XZ -plane. What can you say about its y -coordinate?

Answer :

If a point is in the XZ plane, then its y -coordinate is zero.

Q3 :

Name the octants in which the following points lie:

$(1, 2, 3)$, $(4, -2, 3)$, $(4, -2, -5)$, $(4, 2, -5)$, $(-4, 2, -5)$, $(-4, 2, 5)$, $(-3, -1, 6)$, $(2, -4, -7)$

Answer :

The x -coordinate, y -coordinate, and z -coordinate of point $(1, 2, 3)$ are all positive. Therefore, this point lies in octant **I**.

The x -coordinate, y -coordinate, and z -coordinate of point $(4, -2, 3)$ are positive, negative, and positive respectively. Therefore, this point lies in octant **IV**.

The x -coordinate, y -coordinate, and z -coordinate of point $(4, -2, -5)$ are positive, negative, and negative respectively. Therefore, this point lies in octant **VIII**.

The x -coordinate, y -coordinate, and z -coordinate of point $(4, 2, -5)$ are positive, positive, and negative respectively. Therefore, this point lies in octant **V**.

The x -coordinate, y -coordinate, and z -coordinate of point $(-4, 2, -5)$ are negative, positive, and negative respectively. Therefore, this point lies in octant **VI**.



www.eduinput.com

The x-coordinate, y-coordinate, and z-coordinate of point $(-4, 2, 5)$ are negative, positive, and positive respectively. Therefore, this point lies in octant **II**.

The x-coordinate, y-coordinate, and z-coordinate of point $(-3, -1, 6)$ are negative, negative, and positive respectively. Therefore, this point lies in octant **III**.

The x-coordinate, y-coordinate, and z-coordinate of point $(2, -4, -7)$ are positive, negative, and negative respectively.

Therefore, this point lies in octant **VIII**.

Q4 :

Fill in the blanks:

Answer :

(i) The x-axis and y-axis taken together determine a plane known as

XY – plane

$(x, y, 0)$

(ii) The coordinates of points in the XY-plane are of the form (iii) Coordinate planes divide the space into eight octants.

Exercise 12.2 : Solutions of Questions on Page Number : 273

Q1 :

Find the distance between the following pairs of points:

(i) $(2, 3, 5)$ and $(4, 3, 1)$ (ii) $(-3, 7, 2)$ and $(2, 4, -1)$

(iii) $(-1, 3, -4)$ and $(1, -3, 4)$ (iv) $(2, -1, 3)$ and $(-2, 1, 3)$

Answer :

The distance between points $P(x_1, y_1, z_1)$ and $P(x_2, y_2, z_2)$ is given

by $PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$

(i) Distance between points (2, 3, 5) and (4, 3, 1)

$$\begin{aligned} &= \sqrt{(4-2)^2 + (3-3)^2 + (1-5)^2} \\ &= \sqrt{(2)^2 + (0)^2 + (-4)^2} \\ &= \sqrt{4+16} \\ &= \sqrt{20} \\ &= 2\sqrt{5} \end{aligned}$$

(ii) Distance between points (3, 7, 2) and (2, 4, -1)

$$\begin{aligned} &= \sqrt{(2+3)^2 + (4-7)^2 + (-1-2)^2} \\ &= \sqrt{(5)^2 + (-3)^2 + (-3)^2} \\ &= \sqrt{25+9+9} \\ &= \sqrt{43} \end{aligned}$$

(iii) Distance between points (-1, 3, -4) and (1, -3, 4)

$$\begin{aligned} &= \sqrt{(1+1)^2 + (-3-3)^2 + (4+4)^2} \\ &= \sqrt{(2)^2 + (-6)^2 + (8)^2} \\ &= \sqrt{4+36+64} = \sqrt{104} = 2\sqrt{26} \end{aligned}$$

(iv) Distance between points (2, -1, 3) and (-2, 1, 3)

$$\begin{aligned} &= \sqrt{(-2-2)^2 + (1+1)^2 + (3-3)^2} \\ &= \sqrt{(-4)^2 + (2)^2 + (0)^2} \\ &= \sqrt{16+4} \\ &= \sqrt{20} \\ &= 2\sqrt{5} \end{aligned}$$

Q2 :

Show that the points (-2, 3, 5), (1, 2, 3) and (7, 0, -1) are collinear.

Answer :

Let points $(-2, 3, 5)$, $(1, 2, 3)$, and $(7, 0, -1)$ be denoted by P, Q, and R respectively.

Points P, Q, and R are collinear if they lie on a line.

$$\begin{aligned} PQ &= \sqrt{(1+2)^2 + (2-3)^2 + (3-5)^2} \\ &= \sqrt{(3)^2 + (-1)^2 + (-2)^2} \\ &= \sqrt{9+1+4} \\ &= \sqrt{14} \end{aligned}$$

$$\begin{aligned} QR &= \sqrt{(7-1)^2 + (0-2)^2 + (-1-3)^2} \\ &= \sqrt{(6)^2 + (-2)^2 + (-4)^2} \\ &= \sqrt{36+4+16} \\ &= \sqrt{56} \\ &= 2\sqrt{14} \end{aligned}$$

$$\begin{aligned} PR &= \sqrt{(7+2)^2 + (0-3)^2 + (-1-5)^2} \\ &= \sqrt{(9)^2 + (-3)^2 + (-6)^2} \\ &= \sqrt{81+9+36} \\ &= \sqrt{126} \\ &= 3\sqrt{14} \end{aligned}$$

$$\text{Here, } PQ + QR = \sqrt{14} + 2\sqrt{14} = 3\sqrt{14} = PR$$

Hence, points $P(-2, 3, 5)$, $Q(1, 2, 3)$, and $R(7, 0, -1)$ are collinear.

Q3 :

Verify the following:

- (i) $(0, 7, -10)$, $(1, 6, -6)$ and $(4, 9, -6)$ are the vertices of an isosceles triangle.
- (ii) $(0, 7, 10)$, $(-1, 6, 6)$ and $(-4, 9, 6)$ are the vertices of a right angled triangle.
- (iii) $(-1, 2, 1)$, $(1, -2, 5)$, $(4, -7, 8)$ and $(2, -3, 4)$ are the vertices of a parallelogram.

Answer :

- (i) Let points $(0, 7, -10)$, $(1, 6, -6)$, and $(4, 9, -6)$ be denoted by A, B, and C respectively.

$$\begin{aligned}
 AB &= \sqrt{(1-0)^2 + (6-7)^2 + (-6+10)^2} \\
 &= \sqrt{(1)^2 + (-1)^2 + (4)^2} \\
 &= \sqrt{1+1+16} \\
 &= \sqrt{18} \\
 &= 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 BC &= \sqrt{(4-1)^2 + (9-6)^2 + (-6+6)^2} \\
 &= \sqrt{(3)^2 + (3)^2} \\
 &= \sqrt{9+9} = \sqrt{18} = 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 CA &= \sqrt{(0-4)^2 + (7-9)^2 + (-10+6)^2} \\
 &= \sqrt{(-4)^2 + (-2)^2 + (-4)^2} \\
 &= \sqrt{16+4+16} = \sqrt{36} = 6
 \end{aligned}$$

Here, $AB = BC \neq CA$

Thus, the given points are the vertices of an isosceles triangle.

(ii) Let $(0, 7, 10)$, $(-1, 6, 6)$, and $(-4, 9, 6)$ be denoted by A, B, and C respectively.

$$\begin{aligned}
 AB &= \sqrt{(-1-0)^2 + (6-7)^2 + (6-10)^2} \\
 &= \sqrt{(-1)^2 + (-1)^2 + (-4)^2} \\
 &= \sqrt{1+1+16} = \sqrt{18} \\
 &= 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 BC &= \sqrt{(-4+1)^2 + (9-6)^2 + (6-6)^2} \\
 &= \sqrt{(-3)^2 + (3)^2 + (0)^2} \\
 &= \sqrt{9+9} = \sqrt{18} \\
 &= 3\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 CA &= \sqrt{(0+4)^2 + (7-9)^2 + (10-6)^2} \\
 &= \sqrt{(4)^2 + (-2)^2 + (4)^2} \\
 &= \sqrt{16+4+16} \\
 &= \sqrt{36} \\
 &= 6
 \end{aligned}$$

$$\text{Now, } AB^2 + BC^2 = (3\sqrt{2})^2 + (3\sqrt{2})^2 = 18 + 18 = 36 = AC^2$$

Therefore, by Pythagoras theorem, ABC is a right triangle.

Hence, the given points are the vertices of a right-angled triangle.

(iii) Let $(\hat{a}^{\circ}1, 2, 1)$, $(1, \hat{a}^{\circ}2, 5)$, $(4, \hat{a}^{\circ}7, 8)$, and $(2, \hat{a}^{\circ}3, 4)$ be denoted by A, B, C, and D respectively.

$$\begin{aligned}
 AB &= \sqrt{(1+1)^2 + (-2-2)^2 + (5-1)^2} \\
 &= \sqrt{4+16+16} \\
 &= \sqrt{36} \\
 &= 6
 \end{aligned}$$

$$\begin{aligned}
 BC &= \sqrt{(4-1)^2 + (-7+2)^2 + (8-5)^2} \\
 &= \sqrt{9+25+9} = \sqrt{43}
 \end{aligned}$$

$$\begin{aligned}
 CD &= \sqrt{(2-4)^2 + (-3+7)^2 + (4-8)^2} \\
 &= \sqrt{4+16+16} \\
 &= \sqrt{36} \\
 &= 6
 \end{aligned}$$

$$\begin{aligned}
 DA &= \sqrt{(-1-2)^2 + (2+3)^2 + (1-4)^2} \\
 &= \sqrt{9+25+9} = \sqrt{43}
 \end{aligned}$$

Here, $AB = CD = 6$, $BC = AD = \sqrt{43}$

Hence, the opposite sides of quadrilateral ABCD, whose vertices are taken in order, are equal.

Therefore, ABCD is a parallelogram.

Hence, the given points are the vertices of a parallelogram.



www.eduinput.com

Q4 :

Find the equation of the set of points which are equidistant from the points (1, 2, 3) and (3, 2, -1).

Answer :

Let P (x, y, z) be the point that is equidistant from points A(1, 2, 3) and B(3, 2, -1).

Accordingly, PA = PB

$$\begin{aligned} \Rightarrow PA^2 &= PB^2 \\ \Rightarrow (x-1)^2 + (y-2)^2 + (z-3)^2 &= (x-3)^2 + (y-2)^2 + (z+1)^2 \end{aligned}$$

$$\Rightarrow x^2 - 2x + 1 + y^2 - 4y + 4 + z^2 - 6z + 9 = x^2 - 6x + 9 + y^2 - 4y + 4 + z^2 + 2z + 1$$

$$\Rightarrow -2x - 4y - 6z + 14 = -6x - 4y + 2z + 14 \Rightarrow -2x - 6z + 6x - 2z = 0$$

$$\Rightarrow 4x - 8z = 0$$

$$\Rightarrow x - 2z = 0$$

Thus, the required equation is $x - 2z = 0$.

Q5 :

Find the equation of the set of points P, the sum of whose distances from A (4, 0, 0) and B (-4, 0, 0) is equal to 10.

Answer :

Let the coordinates of P be (x, y, z).

The coordinates of points A and B are (4, 0, 0) and (-4, 0, 0) respectively.

It is given that PA + PB = 10.

$$\begin{aligned} \Rightarrow \sqrt{(x-4)^2 + y^2 + z^2} + \sqrt{(x+4)^2 + y^2 + z^2} &= 10 \\ \Rightarrow \sqrt{(x-4)^2 + y^2 + z^2} &= 10 - \sqrt{(x+4)^2 + y^2 + z^2} \end{aligned}$$

On squaring both sides, we obtain

$$\begin{aligned} \Rightarrow (x-4)^2 + y^2 + z^2 &= 100 - 20\sqrt{(x+4)^2 + y^2 + z^2} + (x+4)^2 + y^2 + z^2 \\ \Rightarrow x^2 - 8x + 16 + y^2 + z^2 &= 100 - 20\sqrt{x^2 + 8x + 16 + y^2 + z^2} + x^2 + 8x + 16 + y^2 + z^2 \\ \Rightarrow 20\sqrt{x^2 + 8x + 16 + y^2 + z^2} &= 100 + 16x \\ \Rightarrow 5\sqrt{x^2 + 8x + 16 + y^2 + z^2} &= (25 + 4x) \end{aligned}$$

On squaring both sides again, we obtain

$$25(x^2 + 8x + 16 + y^2 + z^2) = 625 + 16x^2 + 200x$$

$$\Rightarrow 25x^2 + 200x + 400 + 25y^2 + 25z^2 = 625 + 16x^2 + 200x$$

$$\Rightarrow 9x^2 + 25y^2 + 25z^2 - 225 = 0$$

Thus, the required equation is $9x^2 + 25y^2 + 25z^2 - 225 = 0$.

Exercise 12.3 : Solutions of Questions on Page Number : 277

Q1 :

Find the coordinates of the point which divides the line segment joining the points (-2, 3, 5) and (1, -4, 6) in the ratio (i) 2:3 internally, (ii) 2:3 externally.

Answer :

(i) The coordinates of point R that divides the line segment joining points P (x_1, y_1, z_1) and Q (x_2, y_2, z_2) internally in the ratio $m: n$ are

$$\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$$

Let R (x, y, z) be the point that divides the line segment joining points (-2, 3, 5) and (1, -4, 6) internally in the ratio 2:3

$$x = \frac{2(1) + 3(-2)}{2+3}, y = \frac{2(-4) + 3(3)}{2+3}, \text{ and } z = \frac{2(6) + 3(5)}{2+3}$$

$$\text{i.e., } x = \frac{-4}{5}, y = \frac{1}{5}, \text{ and } z = \frac{27}{5}$$

$$\left(-\frac{4}{5}, \frac{1}{5}, \frac{27}{5} \right)$$

Thus, the coordinates of the required point are $\left(-\frac{4}{5}, \frac{1}{5}, \frac{27}{5} \right)$.

(ii) The coordinates of point R that divides the line segment joining points P (x_1, y_1, z_1) and Q (x_2, y_2, z_2) externally in the ratio $m: n$ are

$$\left(\frac{mx_2 - nx_1}{m-n}, \frac{my_2 - ny_1}{m-n}, \frac{mz_2 - nz_1}{m-n} \right)$$

Let R (x, y, z) be the point that divides the line segment joining points (-2, 3, 5) and (1, -4, 6) externally in the ratio 2:3

$$x = \frac{2(1) - 3(-2)}{2-3}, y = \frac{2(-4) - 3(3)}{2-3}, \text{ and } z = \frac{2(6) - 3(5)}{2-3}$$

$$\text{i.e., } x = -8, y = 17, \text{ and } z = 3$$

Thus, the coordinates of the required point are (-8, 17, 3).

Q2 :

Given that P (3, 2, -4), Q (5, 4, -6) and R (9, 8, -10) are collinear. Find the ratio in which Q divides PR.

Answer :

Let point Q (5, 4, -6) divide the line segment joining points P (3, 2, -4) and R (9, 8, -10) in the ratio $k:1$.

Therefore, by section formula,

$$(5, 4, -6) = \left(\frac{k(9)+3}{k+1}, \frac{k(8)+2}{k+1}, \frac{k(-10)-4}{k+1} \right)$$

$$\Rightarrow \frac{9k+3}{k+1} = 5$$

$$\Rightarrow 9k+3 = 5k+5$$

$$\Rightarrow 4k = 2$$

$$\Rightarrow k = \frac{2}{4} = \frac{1}{2}$$

Thus, point Q divides PR in the ratio 1:2.

Q3 :

Find the ratio in which the YZ-plane divides the line segment formed by joining the points (-2, 4, 7) and (3, -5, 8).

Answer :

Let the YZ plane divide the line segment joining points (-2, 4, 7) and (3, -5, 8) in the ratio $k:1$.

Hence, by section formula, the coordinates of point of intersection are given by

$$\left(\frac{k(3)-2}{k+1}, \frac{k(-5)+4}{k+1}, \frac{k(8)+7}{k+1} \right)$$

On the YZ plane, the x-coordinate of any point is zero.

$$\frac{3k-2}{k+1} = 0$$

$$\Rightarrow 3k-2 = 0$$

$$\Rightarrow k = \frac{2}{3}$$

Thus, the YZ plane divides the line segment formed by joining the given points in the ratio 2:3.

Q4 :

Using section formula, show that the points A (2, -3, 4), B (-1, 2, 1) and $C\left(0, \frac{1}{3}, 2\right)$ are collinear.

Answer :

The given points are A (2, -3, 4), B (-1, 2, 1), and $C\left(0, \frac{1}{3}, 2\right)$.

Let P be a point that divides AB in the ratio $k:1$.

Hence, by section formula, the coordinates of P are given by

$$\left(\frac{k(-1)+2}{k+1}, \frac{k(2)-3}{k+1}, \frac{k(1)+4}{k+1}\right)$$

Now, we find the value of k at which point P coincides with point C.

By taking $\frac{-k+2}{k+1} = 0$, we obtain $k = 2$.

For $k = 2$, the coordinates of point P are $\left(0, \frac{1}{3}, 2\right)$.

i.e., $C\left(0, \frac{1}{3}, 2\right)$ is a point that divides AB externally in the ratio 2:1 and is the same as point P.


Hence, points A, B, and C are collinear.

Q5 :

Find the coordinates of the points which trisect the line segment joining the points P (4, 2, -6) and Q (10, -16, 6).

Answer :

Let A and B be the points that trisect the line segment joining points P (4, 2, -6) and Q (10, -16, 6)



Point A divides PQ in the ratio 1:2. Therefore, by section formula, the coordinates of point A are given by

$$\left(\frac{1(10)+2(4)}{1+2}, \frac{1(-16)+2(2)}{1+2}, \frac{1(6)+2(-6)}{1+2}\right) = (6, -4, -2)$$

Point B divides PQ in the ratio 2:1. Therefore, by section formula, the coordinates of point B are given by

$$\left(\frac{2(10)+1(4)}{2+1}, \frac{2(-16)+1(2)}{2+1}, \frac{2(6)-1(6)}{2+1}\right) = (8, -10, 2)$$

Thus, $(6, -4, 2)$ and $(8, -10, 2)$ are the points that trisect the line segment joining points $P(4, 2, -6)$ and $Q(10, -16, 6)$.


Exercise Miscellaneous : Solutions of Questions on Page Number : 278

Q1 :

Three vertices of a parallelogram ABCD are A $(3, -1, 2)$, B $(1, 2, -4)$ and C $(-1, 1, 2)$. Find the coordinates of the fourth vertex.

Answer :

The three vertices of a parallelogram ABCD are given as A $(3, -1, 2)$, B $(1, 2, -4)$, and C $(-1, 1, 2)$. Let the coordinates of the fourth vertex be D (x, y, z) .



We know that the diagonals of a parallelogram bisect each other.

Therefore, in parallelogram ABCD, AC and BD bisect each other.

\therefore Mid-point of AC = Mid-point of BD

$$\begin{aligned} \Rightarrow \left(\frac{3-1}{2}, \frac{-1+1}{2}, \frac{2+2}{2} \right) &= \left(\frac{x+1}{2}, \frac{y+2}{2}, \frac{z-4}{2} \right) \\ \Rightarrow (1, 0, 2) &= \left(\frac{x+1}{2}, \frac{y+2}{2}, \frac{z-4}{2} \right) \\ \Rightarrow \frac{x+1}{2} = 1, \frac{y+2}{2} = 0, \text{ and } \frac{z-4}{2} = 2 \end{aligned}$$

$$\Rightarrow x = 1, y = -2, \text{ and } z = 8$$


Thus, the coordinates of the fourth vertex are $(1, -2, 8)$.

Q2 :

Find the lengths of the medians of the triangle with vertices A $(0, 0, 6)$, B $(0, 4, 0)$ and $(6, 0, 0)$.

Answer :

Let AD, BE, and CF be the medians of the given triangle ABC.



Since AD is the median, D is the mid-point of BC.

$$\therefore \text{Coordinates of point D} = \left(\frac{0+6}{2}, \frac{4+0}{2}, \frac{0+0}{2} \right) = (3, 2, 0)$$

$$AD = \sqrt{(0-3)^2 + (0-2)^2 + (6-0)^2} = \sqrt{9+4+36} = \sqrt{49} = 7$$

Since BE is the median, E is the mid-point of AC.

$$\therefore \text{Coordinates of point E} = \left(\frac{0+6}{2}, \frac{0+0}{2}, \frac{6+0}{2} \right) = (3, 0, 3)$$

$$BE = \sqrt{(3-0)^2 + (0-4)^2 + (3-0)^2} = \sqrt{9+16+9} = \sqrt{34}$$

Since CF is the median, F is the mid-point of AB.

$$\therefore \text{Coordinates of point F} = \left(\frac{0+0}{2}, \frac{0+4}{2}, \frac{6+0}{2} \right) = (0, 2, 3)$$


$$\text{Length of CF} = \sqrt{(6-0)^2 + (0-2)^2 + (0-3)^2} = \sqrt{36+4+9} = \sqrt{49} = 7$$

Thus, the lengths of the medians of ΔABC are $7, \sqrt{34}$, and 7 .

Q3 :

If the origin is the centroid of the triangle PQR with vertices P $(2a, 2, 6)$, Q $(-4, 3b, -10)$ and R $(8, 14, 2c)$, then find the values of a, b and c .

Answer :



It is known that the coordinates of the centroid of the triangle, whose vertices are (x_1, y_1, z_1) , (x_2, y_2, z_2) and (x_3, y_3, z_3) ,

$$\text{are } \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right).$$

Therefore, coordinates of the centroid of

$$\Delta PQR = \left(\frac{2a - 4 + 8}{3}, \frac{2 + 3b + 14}{3}, \frac{6 - 10 + 2c}{3} \right) = \left(\frac{2a + 4}{3}, \frac{3b + 16}{3}, \frac{2c - 4}{3} \right)$$

It is given that origin is the centroid of ΔPQR .

$$\begin{aligned} \therefore (0, 0, 0) &= \left(\frac{2a + 4}{3}, \frac{3b + 16}{3}, \frac{2c - 4}{3} \right) \\ \Rightarrow \frac{2a + 4}{3} = 0, \frac{3b + 16}{3} = 0 \text{ and } \frac{2c - 4}{3} = 0 \\ \Rightarrow a = -2, b = -\frac{16}{3} \text{ and } c = 2 \end{aligned}$$

$$-2, -\frac{16}{3}, \text{ and } 2.$$

Thus, the respective values of a , b , and c are

Q4 :

Find the coordinates of a point on y -axis which are at a distance of $5\sqrt{2}$ from the point P (3, $\hat{\epsilon}$ 2, 5).

Answer :

If a point is on the y -axis, then x -coordinate and the z -coordinate of the point are zero.

Let A (0, b , 0) be the point on the y -axis at a distance of $5\sqrt{2}$ from point P (3, $\hat{\epsilon}$ 2, 5). Accordingly, $AP = 5\sqrt{2}$

$$\therefore AP^2 = 50$$

$$\Rightarrow (3-0)^2 + (-2-b)^2 + (5-0)^2 = 50$$

$$\Rightarrow 9 + 4 + b^2 + 4b + 25 = 50$$

$$\Rightarrow b^2 + 4b - 12 = 0$$

$$\Rightarrow b^2 + 6b - 2b - 12 = 0$$

$$\Rightarrow (b+6)(b-2) = 0$$

$$\Rightarrow b = -6 \text{ or } 2$$

Thus, the coordinates of the required points are (0, 2, 0) and (0, -6, 0).

Q5 :

A point R with x-coordinate 4 lies on the line segment joining the points P (2, -3, 4) and Q (8, 0, 10). Find the coordinates of the point R.

[Hint suppose R divides PQ in the ratio k: 1. The coordinates of the point R are given by

$$\left(\frac{8k+2}{k+1}, \frac{-3}{k+1}, \frac{10k+4}{k+1} \right)]$$

Answer :

The coordinates of points P and Q are given as P (2, -3, 4) and Q (8, 0, 10).

Let R divide line segment PQ in the ratio k:1.

Hence, by section formula, the coordinates of point R are given by

$$\left(\frac{k(8)+2}{k+1}, \frac{k(0)-3}{k+1}, \frac{k(10)+4}{k+1} \right) = \left(\frac{8k+2}{k+1}, \frac{-3}{k+1}, \frac{10k+4}{k+1} \right)$$

It is given that the x-coordinate of point R is 4.

$$\therefore \frac{8k+2}{k+1} = 4$$

$$\Rightarrow 8k+2 = 4k+4$$

$$\Rightarrow 4k = 2$$

$$\Rightarrow k = \frac{1}{2}$$

$$\left(4, \frac{-3}{\frac{1}{2}+1}, \frac{10\left(\frac{1}{2}\right)+4}{\frac{1}{2}+1} \right) = (4, -2, 6)$$

Therefore, the coordinates of point R are

Q6 :

If A and B be the points (3, 4, 5) and (-1, 3, -7), respectively, find the equation of the set of points P such that $PA^2 + PB^2 = k^2$, where k is a constant.

Answer :

The coordinates of points A and B are given as (3, 4, 5) and (-1, 3, -7) respectively.

Let the coordinates of point P be (x, y, z).

On using distance formula, we obtain

$$\begin{aligned}PA^2 &= (x-3)^2 + (y-4)^2 + (z-5)^2 \\ &= x^2 + 9 - 6x + y^2 + 16 - 8y + z^2 + 25 - 10z \\ &= x^2 - 6x + y^2 - 8y + z^2 - 10z + 50 \\ PB^2 &= (x+1)^2 + (y-3)^2 + (z+7)^2 \\ &= x^2 + 2x + y^2 - 6y + z^2 + 14z + 59\end{aligned}$$

Now, if $PA^2 + PB^2 = k^2$, then

$$\begin{aligned}(x^2 - 6x + y^2 - 8y + z^2 - 10z + 50) + (x^2 + 2x + y^2 - 6y + z^2 + 14z + 59) &= k^2 \\ \Rightarrow 2x^2 + 2y^2 + 2z^2 - 4x - 14y + 4z + 109 &= k^2 \\ \Rightarrow 2(x^2 + y^2 + z^2 - 2x - 7y + 2z) &= k^2 - 109 \\ \Rightarrow x^2 + y^2 + z^2 - 2x - 7y + 2z &= \frac{k^2 - 109}{2}\end{aligned}$$

Thus, the required equation is $x^2 + y^2 + z^2 - 2x - 7y + 2z = \frac{k^2 - 109}{2}$.