



# Velocity | Definition Types and Examples

---

There is a difference between speed and velocity. As distance and displacement are two different quantities, the concept of speed and velocity is also different. Speed depends on distance and velocity depends on displacement. The motion of the body can be described by the [velocity-time graph](#).

## Speed

***The rate of change in the distance of a body is called speed.***

Speed is a scalar quantity. Its unit is a meter per second (m/s).

## Velocity

***The rate of change of displacement is called velocity.***

It is a vector quantity and its direction is along the direction of displacement.

The SI unit of velocity is meter per sec( $\text{ms}^{-1}$ ), having the dimension of  $[\text{LT}^{-1}]$ . The magnitude of velocity is called speed

## Average velocity

***The ratio between the total displacement and the total time taken by the body is called the average velocity.***

If  $d$  is the total displacement and  $t$  is the total time then

$$\text{Average velocity} = \frac{\text{total displacement}}{\text{total time}}$$
$$V_{av} = \frac{d}{t}$$

The average velocity does not tell about

- The actual motion between any two points gives information only about the total path.
- What the path was i.e whether it was straight or curved.
- The nature of the motion that whether it was steady or variable.


A squash ball comes back to its starting point after bouncing off the wall a number of times. Its total displacement is zero so its average velocity is also zero.

## Instantaneous velocity

***The velocity of the body at any instant of time, or at any point of its path with the limit of that time interval tends to zero.***

The limiting value of  $\Delta d / \Delta t$  as time interval  $\Delta t$  approaches zero.

In order to understand the instantaneous velocity, consider a body moving along a path ABC in the XY plane as shown in the adjacent figure.



At any time  $t$ , let the body be at point A. Its position is given by position vector  $r_1$ . After a short interval of time  $\Delta t$ , the body reaches at point B which is given by the position vector  $r_2$ .

The displacement of the body during this short interval of time is given by

$$\Delta d = r_2 - r_1$$

The instantaneous velocity at point A can be found by taking the time interval  $\Delta t$  smaller and smaller.

Then the displacement  $\Delta d$  will become smaller and smaller and ultimately approaches 0.

Thus the velocity at a particular position along the whole path can be found.

During this time term,  $\Delta d/\Delta t$  does not completely vanish to zero. The direction of instantaneous velocity at A is along the tangent at A.

The average velocity of the body may be zero even though its instantaneous velocity is not zero.

## Uniform velocity

***If the instantaneous velocity doesn't change body is said to be moving with uniform velocity.***

If the average and the instantaneous velocities of the moving object are equal it is said to be moving with uniform velocity.

## Variable velocity

***If the instantaneous velocity changes, then a body is said to be moving with variable velocity.***

or if a body covers unequal displacement in an equal interval of time then the body is moving with variable velocity.