

## NCERT Solutions for Class 11 Maths Chapter 3

### Trigonometric Functions Class 11

Chapter 3 Trigonometric Functions Exercise 3.1, 3.2, 3.3, 3.4, miscellaneous Solutions

**Exercise 3.1 : Solutions of Questions on Page Number : 54**

**Q1 :**

**Find the radian measures corresponding to the following degree measures:**

(i)  $25^\circ$  (ii)  $-47^\circ 30'$  (iii)  $240^\circ$  (iv)  $520^\circ$

**Answer :**

(i)  $25^\circ$

We know that  $180^\circ = \pi$  radian

$$\therefore 25^\circ = \frac{\pi}{180} \times 25 \text{ radian} = \frac{5\pi}{36} \text{ radian}$$

(ii)  $-47^\circ 30'$

$$\begin{aligned} -47^\circ 30' &= -47\frac{1}{2} \text{ degree } [1^\circ = 60'] \\ &= \frac{-95}{2} \text{ degree} \end{aligned}$$

Since  $180^\circ = \pi$  radian

$$\begin{aligned} \frac{-95}{2} \text{ degree} &= \frac{\pi}{180} \times \left(\frac{-95}{2}\right) \text{ radian} = \left(\frac{-19}{36 \times 2}\right) \pi \text{ radian} = \frac{-19}{72} \pi \text{ radian} \\ \therefore -47^\circ 30' &= \frac{-19}{72} \pi \text{ radian} \end{aligned}$$

(iii)  $240^\circ$

We know that  $180^\circ = \pi$  radian

$$\therefore 240^\circ = \frac{\pi}{180} \times 240 \text{ radian} = \frac{4}{3} \pi \text{ radian}$$

(iv)  $520^\circ$

We know that  $180^\circ = \pi$  radian

$$\therefore 520^\circ = \frac{\pi}{180} \times 520 \text{ radian} = \frac{26\pi}{9} \text{ radian}$$

**Q2 :**

Find the degree measures corresponding to the following radian measures

$\left( \text{Use } \pi = \frac{22}{7} \right)$

(i)  $\frac{11}{16}$  (ii)  $\frac{5\pi}{3}$  (iii)  $\frac{7\pi}{6}$

**Answer :**

(i)  $\frac{11}{16}$

We know that  $\pi$  radian =  $180^\circ$

$$\begin{aligned} \therefore \frac{11}{16} \text{ radian} &= \frac{180}{\pi} \times \frac{11}{16} \text{ deg ree} = \frac{45 \times 11}{\pi \times 4} \text{ deg ree} \\ &= \frac{45 \times 11 \times 7}{22 \times 4} \text{ deg ree} = \frac{315}{8} \text{ deg ree} \\ &= 39 \frac{3}{8} \text{ deg ree} \\ &= 39^\circ + \frac{3 \times 60}{8} \text{ min utes} \quad [1^\circ = 60'] \\ &= 39^\circ + 22' + \frac{1}{2} \text{ min utes} \\ &= 39^\circ 22' 30'' \quad [1' = 60''] \end{aligned}$$

(ii)  $\frac{5\pi}{3}$

We know that  $\pi$  radian =  $180^\circ$

$$\begin{aligned}
 -4 \text{ radian} &= \frac{180}{\pi} \times (-4) \text{ deg ree} = \frac{180 \times 7(-4)}{22} \text{ deg ree} \\
 &= \frac{-2520}{11} \text{ deg ree} = -229 \frac{1}{11} \text{ deg ree} \\
 &= -229^\circ + \frac{1 \times 60}{11} \text{ min utes} \quad [1^\circ = 60'] \\
 &= -229^\circ + 5' + \frac{5}{11} \text{ min utes} \\
 &= -229^\circ 5' 27'' \quad [1' = 60'']
 \end{aligned}$$

(iii)  $\frac{5\pi}{3}$

We know that  $\pi$  radian =  $180^\circ$

$$\therefore \frac{5\pi}{3} \text{ radian} = \frac{180}{\pi} \times \frac{5\pi}{3} \text{ deg ree} = 300^\circ$$

(iv)  $\frac{7\pi}{6}$

We know that  $\pi$  radian =  $180^\circ$

$$\therefore \frac{7\pi}{6} \text{ radian} = \frac{180}{\pi} \times \frac{7\pi}{6} = 210^\circ$$

**Q3 :**

**A wheel makes 360 revolutions in one minute. Through how many radians does it turn in one second?**

**Answer :**

Number of revolutions made by the wheel in 1 minute = 360

$$\therefore \text{Number of revolutions made by the wheel in 1 second} = \frac{360}{60} = 6$$

In one complete revolution, the wheel turns an angle of  $2\pi$  radian.

Hence, in 6 complete revolutions, it will turn an angle of  $6 \times 2\pi$  radian, i.e.,

$12\pi$  radian

Thus, in one second, the wheel turns an angle of  $12\pi$  radian.

**Q4 :**

Find the degree measure of the angle subtended at the centre of a circle of radius 100 cm by an arc of length

22 cm  $\left( \text{Use } \pi = \frac{22}{7} \right)$ .

**Answer :**

We know that in a circle of radius  $r$  unit, if an arc of length  $l$  unit subtends an angle  $\theta$  radian at the centre, then

Thus, the required angle is  $12^\circ 36''$ .

**Q5 :**


In a circle of diameter 40 cm, the length of a chord is 20 cm. Find the length of minor arc of the chord.

**Answer :**

Diameter of the circle = 40 cm

$$\therefore \text{Radius } (r) \text{ of the circle} = \frac{40}{2} \text{ cm} = 20 \text{ cm}$$

Let AB be a chord (length = 20 cm) of the circle.



In  $\triangle OAB$ ,  $OA = OB = \text{Radius of circle} = 20 \text{ cm}$

Also,  $AB = 20 \text{ cm}$

Thus,  $\triangle OAB$  is an equilateral triangle.

$$\theta = \frac{l}{r}$$

Therefore, for  $r = 100 \text{ cm}$ ,  $l = 22 \text{ cm}$ , we have

$$\begin{aligned} \theta &= \frac{22}{100} \text{ radian} = \frac{180}{\pi} \times \frac{22}{100} \text{ deg ree} = \frac{180 \times 7 \times 22}{22 \times 100} \text{ deg ree} \\ &= \frac{126}{10} \text{ deg ree} = 12\frac{3}{5} \text{ deg ree} = 12^\circ 36' \quad [1^\circ = 60'] \end{aligned}$$

$$\therefore \theta = 60^\circ = \frac{\pi}{3} \text{ radian}$$

We know that in a circle of radius  $r$  unit, if an arc of length  $l$  unit subtends an angle  $\theta$  radian at the centre, then  $\theta = \frac{l}{r}$

$$\frac{\pi}{3} = \frac{\widehat{AB}}{20} \Rightarrow \widehat{AB} = \frac{20\pi}{3} \text{ cm}$$

Thus, the length of the minor arc of the chord is  $\frac{20\pi}{3} \text{ cm}$ .

**Q6 :**

**If in two circles, arcs of the same length subtend angles  $60^\circ$  and  $75^\circ$  at the centre, find the ratio of their radii.**

**Answer :**

$$\text{Now, } 60^\circ = \frac{\pi}{3} \text{ radian}$$

Let the radii of the two circles be  $r_1$  and  $r_2$ . Let an arc of length  $l$  subtend an angle of  $60^\circ$  at the centre of the circle of radius  $r_1$ , while let an arc of length  $l$  subtend an angle of  $75^\circ$  at the centre of the circle of radius  $r_2$ .

$$\theta = \frac{l}{r} \text{ or } l = r\theta$$

$$\text{and } 75^\circ = \frac{5\pi}{12} \text{ radian}$$

We know that in a circle of radius  $r$  unit, if an arc of length  $l$  unit subtends an angle  $\theta$  radian at the centre, then

$$\therefore l = \frac{r_1\pi}{3} \text{ and } l = \frac{r_2 5\pi}{12}$$

$$\Rightarrow \frac{r_1\pi}{3} = \frac{r_2 5\pi}{12}$$

$$\Rightarrow r_1 = \frac{r_2 5}{4}$$

$$\Rightarrow \frac{r_1}{r_2} = \frac{5}{4}$$

Thus, the ratio of the radii is 5:4.

**Q7 :**

Find the angle in radian through which a pendulum swings if its length is 75 cm and the tip describes an arc of length

(i) 10 cm (ii) 15 cm (iii) 21 cm

**Answer :**

$$\theta = \frac{l}{r}$$

We know that in a circle of radius  $r$  unit, if an arc of length  $l$  unit subtends an angle  $\theta$  radian at the centre, then

It is given that  $r = 75$  cm

(i) Here,  $l = 10$  cm

$$\theta = \frac{10}{75} \text{ radian} = \frac{2}{15} \text{ radian}$$

(ii) Here,  $l = 15$  cm

$$\theta = \frac{15}{75} \text{ radian} = \frac{1}{5} \text{ radian}$$

(iii) Here,  $l = 21$  cm

$$\theta = \frac{21}{75} \text{ radian} = \frac{7}{25} \text{ radian}$$

**Exercise 3.2 : Solutions of Questions on Page Number : 63**

**Q1 :**

$$\cos x = -\frac{1}{2}, x \text{ lies in third quadrant.}$$

Find the values of other five trigonometric functions if

Answer :

$$\cos x = -\frac{1}{2}$$

$$\therefore \sec x = \frac{1}{\cos x} = \frac{1}{\left(-\frac{1}{2}\right)} = -2$$

$$\sin^2 x + \cos^2 x = 1$$

$$\Rightarrow \sin^2 x = 1 - \cos^2 x$$

$$\Rightarrow \sin^2 x = 1 - \left(-\frac{1}{2}\right)^2$$

$$\Rightarrow \sin^2 x = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\Rightarrow \sin x = \pm \frac{\sqrt{3}}{2}$$

Since  $x$  lies in the 3<sup>rd</sup> quadrant, the value of  $\sin x$  will be negative.

$$\therefore \sin x = -\frac{\sqrt{3}}{2}$$

$$\operatorname{cosec} x = \frac{1}{\sin x} = \frac{1}{\left(-\frac{\sqrt{3}}{2}\right)} = -\frac{2}{\sqrt{3}}$$

$$\tan x = \frac{\sin x}{\cos x} = \frac{\left(-\frac{\sqrt{3}}{2}\right)}{\left(-\frac{1}{2}\right)} = \sqrt{3}$$

$$\cot x = \frac{1}{\tan x} = \frac{1}{\sqrt{3}}$$

Q2 :

Find the values of other five trigonometric functions if  $\sin x = \frac{3}{5}$ ,  $x$  lies in second quadrant.

Answer :

$$\sin x = \frac{3}{5}$$

$$\operatorname{cosec} x = \frac{1}{\sin x} = \frac{1}{\left(\frac{3}{5}\right)} = \frac{5}{3}$$

$$\sin^2 x + \cos^2 x = 1$$

$$\Rightarrow \cos^2 x = 1 - \sin^2 x$$

$$\Rightarrow \cos^2 x = 1 - \left(\frac{3}{5}\right)^2$$

$$\Rightarrow \cos^2 x = 1 - \frac{9}{25}$$

$$\Rightarrow \cos^2 x = \frac{16}{25}$$

$$\Rightarrow \cos x = \pm \frac{4}{5}$$

Since  $x$  lies in the 2<sup>nd</sup> quadrant, the value of  $\cos x$  will be negative

$$\therefore \cos x = -\frac{4}{5}$$

$$\sec x = \frac{1}{\cos x} = \frac{1}{\left(-\frac{4}{5}\right)} = -\frac{5}{4}$$

$$\tan x = \frac{\sin x}{\cos x} = \frac{\left(\frac{3}{5}\right)}{\left(-\frac{4}{5}\right)} = -\frac{3}{4}$$

$$\cot x = \frac{1}{\tan x} = -\frac{4}{3}$$

Q3 :

Find the values of other five trigonometric functions if  $\cot x = \frac{3}{4}$ ,  $x$  lies in third quadrant.



Answer :

$$\cot x = \frac{3}{4}$$

$$\tan x = \frac{1}{\cot x} = \frac{1}{\left(\frac{3}{4}\right)} = \frac{4}{3}$$

$$1 + \tan^2 x = \sec^2 x$$

$$\Rightarrow 1 + \left(\frac{4}{3}\right)^2 = \sec^2 x$$

$$\Rightarrow 1 + \frac{16}{9} = \sec^2 x$$

$$\Rightarrow \frac{25}{9} = \sec^2 x$$

$$\Rightarrow \sec x = \pm \frac{5}{3}$$

Since  $x$  lies in the 3<sup>rd</sup> quadrant, the value of  $\sec x$  will be negative.

$$\therefore \sec x = -\frac{5}{3}$$

$$\cos x = \frac{1}{\sec x} = \frac{1}{\left(-\frac{5}{3}\right)} = -\frac{3}{5}$$

$$\tan x = \frac{\sin x}{\cos x}$$

$$\Rightarrow \frac{4}{3} = \frac{\sin x}{\left(-\frac{3}{5}\right)}$$

$$\Rightarrow \sin x = \left(\frac{4}{3}\right) \times \left(-\frac{3}{5}\right) = -\frac{4}{5}$$

$$\operatorname{cosec} x = \frac{1}{\sin x} = -\frac{5}{4}$$

Q4 :

Find the values of other five trigonometric functions if  $\sec x = \frac{13}{5}$ ,  $x$  lies in fourth quadrant.

**Answer :**

$$\sec x = \frac{13}{5}$$

$$\cos x = \frac{1}{\sec x} = \frac{1}{\left(\frac{13}{5}\right)} = \frac{5}{13}$$

$$\sin^2 x + \cos^2 x = 1$$

$$\Rightarrow \sin^2 x = 1 - \cos^2 x$$

$$\Rightarrow \sin^2 x = 1 - \left(\frac{5}{13}\right)^2$$

$$\Rightarrow \sin^2 x = 1 - \frac{25}{169} = \frac{144}{169}$$

$$\Rightarrow \sin x = \pm \frac{12}{13}$$

Since  $x$  lies in the 4<sup>th</sup> quadrant, the value of  $\sin x$  will be negative.

$$\therefore \sin x = -\frac{12}{13}$$

$$\operatorname{cosec} x = \frac{1}{\sin x} = \frac{1}{\left(-\frac{12}{13}\right)} = -\frac{13}{12}$$

$$\tan x = \frac{\sin x}{\cos x} = \frac{\left(-\frac{12}{13}\right)}{\left(\frac{5}{13}\right)} = -\frac{12}{5}$$

$$\cot x = \frac{1}{\tan x} = \frac{1}{\left(-\frac{12}{5}\right)} = -\frac{5}{12}$$

**Q5 :**

Find the values of other five trigonometric functions if  $\tan x = -\frac{5}{12}$ ,  $x$  lies in second quadrant.

**Answer :**

$$\tan x = -\frac{5}{12}$$

$$\cot x = \frac{1}{\tan x} = \frac{1}{\left(-\frac{5}{12}\right)} = -\frac{12}{5}$$

$$1 + \tan^2 x = \sec^2 x$$

$$\Rightarrow 1 + \left(-\frac{5}{12}\right)^2 = \sec^2 x$$

$$\Rightarrow 1 + \frac{25}{144} = \sec^2 x$$

$$\Rightarrow \frac{169}{144} = \sec^2 x$$

$$\Rightarrow \sec x = \pm \frac{13}{12}$$

Since  $x$  lies in the 2<sup>nd</sup> quadrant, the value of  $\sec x$  will be negative.

$$\therefore \sec x = -\frac{13}{12}$$

$$\cos x = \frac{1}{\sec x} = \frac{1}{\left(-\frac{13}{12}\right)} = -\frac{12}{13}$$

$$\tan x = \frac{\sin x}{\cos x}$$

$$\Rightarrow -\frac{5}{12} = \frac{\sin x}{\left(-\frac{12}{13}\right)}$$

$$\Rightarrow \sin x = \left(-\frac{5}{12}\right) \times \left(-\frac{12}{13}\right) = \frac{5}{13}$$

$$\operatorname{cosec} x = \frac{1}{\sin x} = \frac{1}{\left(\frac{5}{13}\right)} = \frac{13}{5}$$

**Q6 :**

Find the value of the trigonometric function  $\sin 765^\circ$

**Answer :**

It is known that the values of  $\sin x$  repeat after an interval of  $2\pi$  or  $360^\circ$ .

$$\therefore \sin 765^\circ = \sin (2 \times 360^\circ + 45^\circ) = \sin 45^\circ = \frac{1}{\sqrt{2}}$$

**Q7 :**

**Find the value of the trigonometric function cosec (-1410°)**

**Answer :**

It is known that the values of cosec x repeat after an interval of  $2\pi$  or  $360^\circ$ .

$$\begin{aligned}\therefore \operatorname{cosec}(-1410^\circ) &= \operatorname{cosec}(-1410^\circ + 4 \times 360^\circ) \\ &= \operatorname{cosec}(-1410^\circ + 1440^\circ) \\ &= \operatorname{cosec}30^\circ = 2\end{aligned}$$

**Q8 :**

**Find the value of the trigonometric function**  $\tan \frac{19\pi}{3}$

**Answer :**

It is known that the values of tan x repeat after an interval of  $\pi$  or  $180^\circ$ .

$$\therefore \tan \frac{19\pi}{3} = \tan 6\frac{1}{3}\pi = \tan \left( 6\pi + \frac{\pi}{3} \right) = \tan \frac{\pi}{3} = \tan 60^\circ = \sqrt{3}$$

**Q9 :**

**Find the value of the trigonometric function**  $\sin \left( -\frac{11\pi}{3} \right)$

**Answer :**

It is known that the values of sin x repeat after an interval of  $2\pi$  or  $360^\circ$ .

$$\therefore \sin \left( -\frac{11\pi}{3} \right) = \sin \left( -\frac{11\pi}{3} + 2 \times 2\pi \right) = \sin \left( \frac{\pi}{3} \right) = \frac{\sqrt{3}}{2}$$

**Q10 :**

**Find the value of the trigonometric function**  $\cot \left( -\frac{15\pi}{4} \right)$

**Answer :**

It is known that the values of  $\cot x$  repeat after an interval of  $\pi$  or  $180^\circ$ .

$$\therefore \cot\left(-\frac{15\pi}{4}\right) = \cot\left(-\frac{15\pi}{4} + 4\pi\right) = \cot\frac{\pi}{4} = 1$$

**Exercise 3.3 :** Solutions of Questions on Page Number : 73

**Q1 :**

$$\sin^2 \frac{\pi}{6} + \cos^2 \frac{\pi}{3} - \tan^2 \frac{\pi}{4} = -\frac{1}{2}$$

**Answer :**

$$\begin{aligned} \text{L.H.S.} &= \sin^2 \frac{\pi}{6} + \cos^2 \frac{\pi}{3} - \tan^2 \frac{\pi}{4} \\ &= \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 - (1)^2 \\ &= \frac{1}{4} + \frac{1}{4} - 1 = -\frac{1}{2} \\ &= \text{R.H.S.} \end{aligned}$$

**Q2 :**

$$\text{Prove that } 2 \sin^2 \frac{\pi}{6} + \operatorname{cosec}^2 \frac{7\pi}{6} \cos^2 \frac{\pi}{3} = \frac{3}{2}$$

**Answer :**

$$\begin{aligned} \text{L.H.S.} &= 2 \sin^2 \frac{\pi}{6} + \operatorname{cosec}^2 \frac{7\pi}{6} \cos^2 \frac{\pi}{3} \\ &= 2 \left(\frac{1}{2}\right)^2 + \operatorname{cosec}^2 \left(\pi + \frac{\pi}{6}\right) \left(\frac{1}{2}\right)^2 \\ &= 2 \times \frac{1}{4} + \left(-\operatorname{cosec} \frac{\pi}{6}\right)^2 \left(\frac{1}{4}\right) \\ &= \frac{1}{2} + (-2)^2 \left(\frac{1}{4}\right) \\ &= \frac{1}{2} + \frac{4}{4} = \frac{1}{2} + 1 = \frac{3}{2} \\ &= \text{R.H.S.} \end{aligned}$$

**Q3 :**

Prove that  $\cot^2 \frac{\pi}{6} + \operatorname{cosec} \frac{5\pi}{6} + 3 \tan^2 \frac{\pi}{6} = 6$

Answer :

$$\text{L.H.S.} = \cot^2 \frac{\pi}{6} + \operatorname{cosec} \frac{5\pi}{6} + 3 \tan^2 \frac{\pi}{6}$$

$$= (\sqrt{3})^2 + \operatorname{cosec} \left( \pi - \frac{\pi}{6} \right) + 3 \left( \frac{1}{\sqrt{3}} \right)^2$$

$$= 3 + \operatorname{cosec} \frac{\pi}{6} + 3 \times \frac{1}{3}$$

$$= 3 + 2 + 1 = 6$$

$$= \text{R.H.S}$$

Q4 :

Prove that  $2 \sin^2 \frac{3\pi}{4} + 2 \cos^2 \frac{\pi}{4} + 2 \sec^2 \frac{\pi}{3} = 10$

Answer :

$$\text{L.H.S} = 2 \sin^2 \frac{3\pi}{4} + 2 \cos^2 \frac{\pi}{4} + 2 \sec^2 \frac{\pi}{3}$$

$$= 2 \left\{ \sin \left( \pi - \frac{\pi}{4} \right) \right\}^2 + 2 \left( \frac{1}{\sqrt{2}} \right)^2 + 2(2)^2$$

$$= 2 \left\{ \sin \frac{\pi}{4} \right\}^2 + 2 \times \frac{1}{2} + 8$$

$$= 2 \left( \frac{1}{\sqrt{2}} \right)^2 + 1 + 8$$

$$= 1 + 1 + 8$$

$$= 10$$

$$= \text{R.H.S}$$

Q5 :

Find the value of:

(i)  $\sin 75^\circ$

(ii)  $\tan 15^\circ$

**Answer :**

$$(i) \sin 75^\circ = \sin (45^\circ + 30^\circ)$$

$$= \sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ$$

$$[\sin (x + y) = \sin x \cos y + \cos x \sin y]$$

$$\begin{aligned} &= \left(\frac{1}{\sqrt{2}}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(\frac{1}{\sqrt{2}}\right)\left(\frac{1}{2}\right) \\ &= \frac{\sqrt{3}}{2\sqrt{2}} + \frac{1}{2\sqrt{2}} = \frac{\sqrt{3}+1}{2\sqrt{2}} \end{aligned}$$

$$(ii) \tan 15^\circ = \tan (45^\circ - 30^\circ)$$

$$= \frac{\tan 45^\circ - \tan 30^\circ}{1 + \tan 45^\circ \tan 30^\circ}$$

$$\left[ \tan (x - y) = \frac{\tan x - \tan y}{1 + \tan x \tan y} \right]$$

$$= \frac{1 - \frac{1}{\sqrt{3}}}{1 + 1\left(\frac{1}{\sqrt{3}}\right)} = \frac{\frac{\sqrt{3}-1}{\sqrt{3}}}{\frac{\sqrt{3}+1}{\sqrt{3}}}$$

$$= \frac{\sqrt{3}-1}{\sqrt{3}+1} = \frac{(\sqrt{3}-1)^2}{(\sqrt{3}+1)(\sqrt{3}-1)} = \frac{3+1-2\sqrt{3}}{(\sqrt{3})^2 - (1)^2}$$

$$= \frac{4-2\sqrt{3}}{3-1} = 2 - \sqrt{3}$$

**Q6 :**

**Prove that:**  $\cos\left(\frac{\pi}{4} - x\right)\cos\left(\frac{\pi}{4} - y\right) - \sin\left(\frac{\pi}{4} - x\right)\sin\left(\frac{\pi}{4} - y\right) = \sin(x + y)$

**Answer :**

$$\begin{aligned}
 & \cos\left(\frac{\pi}{4}-x\right)\cos\left(\frac{\pi}{4}-y\right)-\sin\left(\frac{\pi}{4}-x\right)\sin\left(\frac{\pi}{4}-y\right) \\
 &= \frac{1}{2}\left[2\cos\left(\frac{\pi}{4}-x\right)\cos\left(\frac{\pi}{4}-y\right)\right]+\frac{1}{2}\left[-2\sin\left(\frac{\pi}{4}-x\right)\sin\left(\frac{\pi}{4}-y\right)\right] \\
 &= \frac{1}{2}\left[\cos\left\{\left(\frac{\pi}{4}-x\right)+\left(\frac{\pi}{4}-y\right)\right\}+\cos\left\{\left(\frac{\pi}{4}-x\right)-\left(\frac{\pi}{4}-y\right)\right\}\right] \\
 &+ \frac{1}{2}\left[\cos\left\{\left(\frac{\pi}{4}-x\right)+\left(\frac{\pi}{4}-y\right)\right\}-\cos\left\{\left(\frac{\pi}{4}-x\right)-\left(\frac{\pi}{4}-y\right)\right\}\right] \\
 &\left[\because 2\cos A\cos B=\cos(A+B)+\cos(A-B)\right] \\
 &\left[-2\sin A\sin B=\cos(A+B)-\cos(A-B)\right] \\
 &= 2\times\frac{1}{2}\left[\cos\left\{\left(\frac{\pi}{4}-x\right)+\left(\frac{\pi}{4}-y\right)\right\}\right] \\
 &= \cos\left[\frac{\pi}{2}-(x+y)\right] \\
 &= \sin(x+y) \\
 &= \text{R.H.S}
 \end{aligned}$$

**Q7 :**

$$\frac{\tan\left(\frac{\pi}{4}+x\right)}{\tan\left(\frac{\pi}{4}-x\right)}=\left(\frac{1+\tan x}{1-\tan x}\right)^2$$

**Prove that:**

**Answer :**

$$\tan(A+B)=\frac{\tan A+\tan B}{1-\tan A\tan B} \text{ and } \tan(A-B)=\frac{\tan A-\tan B}{1+\tan A\tan B}$$

It is known that

$$\begin{aligned}
 \therefore \text{L.H.S.} &= \frac{\tan\left(\frac{\pi}{4}+x\right)}{\tan\left(\frac{\pi}{4}-x\right)} = \frac{\left(\frac{\tan\frac{\pi}{4}+\tan x}{1-\tan\frac{\pi}{4}\tan x}\right)}{\left(\frac{\tan\frac{\pi}{4}-\tan x}{1+\tan\frac{\pi}{4}\tan x}\right)} = \frac{\left(\frac{1+\tan x}{1-\tan x}\right)}{\left(\frac{1-\tan x}{1+\tan x}\right)} = \left(\frac{1+\tan x}{1-\tan x}\right)^2 = \text{R.H.S.}
 \end{aligned}$$



**Q8 :**

$$\frac{\cos(\pi + x) \cos(-x)}{\sin(\pi - x) \cos\left(\frac{\pi}{2} + x\right)} = \cot^2 x$$

**Prove that**

**Answer :**

$$\begin{aligned}\text{L.H.S.} &= \frac{\cos(\pi + x) \cos(-x)}{\sin(\pi - x) \cos\left(\frac{\pi}{2} + x\right)} \\ &= \frac{[-\cos x][\cos x]}{(\sin x)(-\sin x)} \\ &= \frac{-\cos^2 x}{-\sin^2 x} \\ &= \cot^2 x \\ &= \text{R.H.S.}\end{aligned}$$

**Q9 :**

$$\cos\left(\frac{3\pi}{2} + x\right) \cos(2\pi + x) \left[ \cot\left(\frac{3\pi}{2} - x\right) + \cot(2\pi + x) \right] = 1$$

**Answer :**

$$\begin{aligned}\text{L.H.S.} &= \cos\left(\frac{3\pi}{2} + x\right) \cos(2\pi + x) \left[ \cot\left(\frac{3\pi}{2} - x\right) + \cot(2\pi + x) \right] \\ &= \sin x \cos x [\tan x + \cot x] \\ &= \sin x \cos x \left( \frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} \right) \\ &= (\sin x \cos x) \left[ \frac{\sin^2 x + \cos^2 x}{\sin x \cos x} \right] \\ &= 1 = \text{R.H.S.}\end{aligned}$$

**Q10 :**

**Prove that  $\sin(n+1)x \sin(n+2)x + \cos(n+1)x \cos(n+2)x = \cos x$**

**Answer :**

$$\text{L.H.S.} = \sin(n+1)x \sin(n+2)x + \cos(n+1)x \cos(n+2)x$$

$$\begin{aligned} &= \frac{1}{2} [2 \sin(n+1)x \sin(n+2)x + 2 \cos(n+1)x \cos(n+2)x] \\ &= \frac{1}{2} \left[ \cos\{(n+1)x - (n+2)x\} - \cos\{(n+1)x + (n+2)x\} \right. \\ &\quad \left. + \cos\{(n+1)x + (n+2)x\} + \cos\{(n+1)x - (n+2)x\} \right] \\ &\quad \left[ \because -2 \sin A \sin B = \cos(A+B) - \cos(A-B) \right. \\ &\quad \left. 2 \cos A \cos B = \cos(A+B) + \cos(A-B) \right] \\ &= \frac{1}{2} \times 2 \cos\{(n+1)x - (n+2)x\} \\ &= \cos(-x) = \cos x = \text{R.H.S.} \end{aligned}$$

**Q11 :**

Prove that  $\cos\left(\frac{3\pi}{4} + x\right) - \cos\left(\frac{3\pi}{4} - x\right) = -\sqrt{2} \sin x$

**Answer :**

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \cdot \sin\left(\frac{A-B}{2}\right)$$

It is known that

$$\therefore \text{L.H.S.} = \cos\left(\frac{3\pi}{4} + x\right) - \cos\left(\frac{3\pi}{4} - x\right)$$

$$= -2 \sin\left\{\frac{\left(\frac{3\pi}{4} + x\right) + \left(\frac{3\pi}{4} - x\right)}{2}\right\} \cdot \sin\left\{\frac{\left(\frac{3\pi}{4} + x\right) - \left(\frac{3\pi}{4} - x\right)}{2}\right\}$$

$$= -2 \sin\left(\frac{3\pi}{4}\right) \sin x$$

$$= -2 \sin\left(\pi - \frac{\pi}{4}\right) \sin x$$

$$= -2 \sin \frac{\pi}{4} \sin x$$

$$= -2 \times \frac{1}{\sqrt{2}} \times \sin x$$

$$= -\sqrt{2} \sin x$$

$$= \text{R.H.S.}$$

**Q12 :**

**Prove that  $\sin^2 6x - \sin^2 4x = \sin 2x \sin 10x$**

**Answer :** It

is known

$$\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right), \quad \sin A - \sin B = 2 \cos\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

that

$$\therefore \text{L.H.S.} = \sin^2 6x - \sin^2 4x$$

$$= (\sin 6x + \sin 4x) (\sin 6x - \sin 4x)$$

$$= \left[ 2 \sin\left(\frac{6x+4x}{2}\right) \cos\left(\frac{6x-4x}{2}\right) \right] \left[ 2 \cos\left(\frac{6x+4x}{2}\right) \sin\left(\frac{6x-4x}{2}\right) \right]$$

$$= (2 \sin 5x \cos x) (2 \cos 5x \sin x) = (2$$

$$\sin 5x \cos 5x) (2 \sin x \cos x)$$

$$= \sin 10x \sin 2x = \text{R.H.S.}$$

**Q13 :**

**Prove that  $\cos^2 2x - \cos^2 6x = \sin 4x \sin 8x$**

**Answer :**

It is known

that 
$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right), \quad \cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

$$\therefore \text{L.H.S.} = \cos^2 2x - \cos^2 6x$$

$$= (\cos 2x + \cos 6x) (\cos 2x - \cos 6x)$$

$$\begin{aligned} &= \left[ 2 \cos\left(\frac{2x+6x}{2}\right) \cos\left(\frac{2x-6x}{2}\right) \right] \left[ -2 \sin\left(\frac{2x+6x}{2}\right) \sin\left(\frac{2x-6x}{2}\right) \right] \\ &= \left[ 2 \cos 4x \cos(-2x) \right] \left[ -2 \sin 4x \sin(-2x) \right] \end{aligned}$$

$$= [2 \cos 4x \cos 2x] [-2 \sin 4x (-\sin 2x)]$$

$$= (2 \sin 4x \cos 4x) (2 \sin 2x \cos 2x)$$

$$= \sin 8x \sin 4x =$$

$$\text{R.H.S.}$$

**Q14 :**

**Prove that  $\sin 2x + 2\sin 4x + \sin 6x = 4\cos^2 x \sin 4x$**

**Answer :**

$$\text{L.H.S.} = \sin 2x + 2 \sin 4x + \sin 6x$$

$$= [\sin 2x + \sin 6x] + 2 \sin 4x$$

$$= \left[ 2 \sin\left(\frac{2x+6x}{2}\right) \cos\left(\frac{2x-6x}{2}\right) \right] + 2 \sin 4x$$

$$\left[ \because \sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) \right]$$

$$= 2 \sin 4x \cos(-2x) + 2 \sin 4x$$

$$= 2 \sin 4x \cos 2x + 2 \sin 4x$$

$$= 2 \sin 4x (\cos 2x + 1)$$

$$= 2 \sin 4x (2 \cos^2 x - 1 + 1)$$

$$= 2 \sin 4x (2 \cos^2 x)$$

$$= 4 \cos^2 x \sin 4x = \text{R.H.S.}$$

**Q15 :**

**Prove that  $\cot 4x (\sin 5x + \sin 3x) = \cot x (\sin 5x - \sin 3x)$**

**Answer :**

$$\text{L.H.S} = \cot 4x (\sin 5x + \sin 3x)$$

$$= \frac{\cos 4x}{\sin 4x} \left[ 2 \sin \left( \frac{5x+3x}{2} \right) \cos \left( \frac{5x-3x}{2} \right) \right]$$

$$\left[ \because \sin A + \sin B = 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right) \right]$$

$$= \left( \frac{\cos 4x}{\sin 4x} \right) [2 \sin 4x \cos x]$$

$$= 2 \cos 4x \cos x$$

$$\text{R.H.S.} = \cot x (\sin 5x - \sin 3x)$$

$$= \frac{\cos x}{\sin x} \left[ 2 \cos \left( \frac{5x+3x}{2} \right) \sin \left( \frac{5x-3x}{2} \right) \right]$$

$$\left[ \because \sin A - \sin B = 2 \cos \left( \frac{A+B}{2} \right) \sin \left( \frac{A-B}{2} \right) \right]$$

$$= \frac{\cos x}{\sin x} [2 \cos 4x \sin x]$$

$$= 2 \cos 4x \cos x$$

$$\text{L.H.S.} = \text{R.H.S.}$$

**Q16 :**

$$\text{Prove that } \frac{\cos 9x - \cos 5x}{\sin 17x - \sin 3x} = -\frac{\sin 2x}{\cos 10x}$$

**Answer :**

It is known that

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right), \quad \sin A - \sin B = 2 \cos\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

$$\therefore \text{L.H.S} = \frac{\cos 9x - \cos 5x}{\sin 17x - \sin 3x}$$

$$\begin{aligned} &= \frac{-2 \sin\left(\frac{9x+5x}{2}\right) \cdot \sin\left(\frac{9x-5x}{2}\right)}{2 \cos\left(\frac{17x+3x}{2}\right) \cdot \sin\left(\frac{17x-3x}{2}\right)} \\ &= \frac{-2 \sin 7x \cdot \sin 2x}{2 \cos 10x \cdot \sin 7x} \\ &= -\frac{\sin 2x}{\cos 10x} \\ &= \text{R.H.S.} \end{aligned}$$

**Q17 :**

**Prove that**  $\frac{\sin 5x + \sin 3x}{\cos 5x + \cos 3x} = \tan 4x$

**Answer :**

It is known that

$$\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right), \quad \cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\therefore \text{L.H.S.} = \frac{\sin 5x + \sin 3x}{\cos 5x + \cos 3x}$$

$$\begin{aligned} &= \frac{2 \sin\left(\frac{5x+3x}{2}\right) \cdot \cos\left(\frac{5x-3x}{2}\right)}{2 \cos\left(\frac{5x+3x}{2}\right) \cdot \cos\left(\frac{5x-3x}{2}\right)} \\ &= \frac{2 \sin 4x \cdot \cos x}{2 \cos 4x \cdot \cos x} \\ &= \frac{\sin 4x}{\cos 4x} \\ &= \tan 4x = \text{R.H.S.} \end{aligned}$$

**Q18 :**

**Prove that**  $\frac{\sin x - \sin y}{\cos x + \cos y} = \tan \frac{x-y}{2}$

**Answer :**

It is known that

$$\sin A - \sin B = 2 \cos \left( \frac{A+B}{2} \right) \sin \left( \frac{A-B}{2} \right), \quad \cos A + \cos B = 2 \cos \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right)$$

$$\therefore \text{L.H.S.} = \frac{\sin x - \sin y}{\cos x + \cos y}$$

$$= \frac{2 \cos \left( \frac{x+y}{2} \right) \cdot \sin \left( \frac{x-y}{2} \right)}{2 \cos \left( \frac{x+y}{2} \right) \cdot \cos \left( \frac{x-y}{2} \right)}$$

$$= \frac{\sin \left( \frac{x-y}{2} \right)}{\cos \left( \frac{x-y}{2} \right)}$$

$$= \tan \left( \frac{x-y}{2} \right) = \text{R.H.S.}$$

**Q19 :**

**Prove that**  $\frac{\sin x + \sin 3x}{\cos x + \cos 3x} = \tan 2x$

**Answer :**

It is known that

$$\sin A + \sin B = 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right), \quad \cos A + \cos B = 2 \cos \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right)$$

$$\therefore \text{L.H.S.} = \frac{\sin x + \sin 3x}{\cos x + \cos 3x}$$

$$\begin{aligned}
 &= \frac{2 \sin\left(\frac{x+3x}{2}\right) \cos\left(\frac{x-3x}{2}\right)}{2 \cos\left(\frac{x+3x}{2}\right) \cos\left(\frac{x-3x}{2}\right)} \\
 &= \frac{\sin 2x}{\cos 2x} \\
 &= \tan 2x \\
 &= \text{R.H.S}
 \end{aligned}$$

**Q20 :**

**Prove that**  $\frac{\sin x - \sin 3x}{\sin^2 x - \cos^2 x} = 2 \sin x$

**Answer :**

It is known that

$$\sin A - \sin B = 2 \cos\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right), \quad \cos^2 A - \sin^2 A = \cos 2A$$

$$\therefore \text{L.H.S.} = \frac{\sin x - \sin 3x}{\sin^2 x - \cos^2 x}$$

$$\begin{aligned}
 &= \frac{2 \cos\left(\frac{x+3x}{2}\right) \sin\left(\frac{x-3x}{2}\right)}{-\cos 2x} \\
 &= \frac{2 \cos 2x \sin(-x)}{-\cos 2x} \\
 &= -2 \times (-\sin x) \\
 &= 2 \sin x = \text{R.H.S.}
 \end{aligned}$$

**Q21 :**

**Prove that**  $\frac{\cos 4x + \cos 3x + \cos 2x}{\sin 4x + \sin 3x + \sin 2x} = \cot 3x$

**Answer :**

$$\text{L.H.S.} = \frac{\cos 4x + \cos 3x + \cos 2x}{\sin 4x + \sin 3x + \sin 2x}$$



$$\begin{aligned}
&= \frac{(\cos 4x + \cos 2x) + \cos 3x}{(\sin 4x + \sin 2x) + \sin 3x} \\
&= \frac{2 \cos \left( \frac{4x+2x}{2} \right) \cos \left( \frac{4x-2x}{2} \right) + \cos 3x}{2 \sin \left( \frac{4x+2x}{2} \right) \cos \left( \frac{4x-2x}{2} \right) + \sin 3x} \\
&\left[ \because \cos A + \cos B = 2 \cos \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right), \sin A + \sin B = 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right) \right] \\
&= \frac{2 \cos 3x \cos x + \cos 3x}{2 \sin 3x \cos x + \sin 3x} \\
&= \frac{\cos 3x (2 \cos x + 1)}{\sin 3x (2 \cos x + 1)} \\
&= \cot 3x = \text{R.H.S.}
\end{aligned}$$

**Q22 :**

**Prove that  $\cot x \cot 2x - \cot 2x \cot 3x - \cot 3x \cot x = 1$**

**Answer :**

$$\text{L.H.S.} = \cot x \cot 2x - \cot 2x \cot 3x - \cot 3x \cot x$$

$$= \cot x \cot 2x - \cot 3x (\cot 2x + \cot x)$$

$$= \cot x \cot 2x - \cot (2x + x) (\cot 2x + \cot x)$$

$$\begin{aligned}
&= \cot x \cot 2x - \left[ \frac{\cot 2x \cot x - 1}{\cot x + \cot 2x} \right] (\cot 2x + \cot x) \\
&\left[ \because \cot(A+B) = \frac{\cot A \cot B - 1}{\cot A + \cot B} \right]
\end{aligned}$$

$$= \cot x \cot 2x - (\cot 2x \cot x - 1) = 1$$

$$= \text{R.H.S.}$$

**Q23 :**

**Prove that** 
$$\tan 4x = \frac{4 \tan x (1 - \tan^2 x)}{1 - 6 \tan^2 x + \tan^4 x}$$

**Answer :**

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

It is known that

$$\therefore \text{L.H.S.} = \tan 4x = \tan 2(2x)$$

$$\begin{aligned} &= \frac{2 \tan 2x}{1 - \tan^2 (2x)} \\ &= \frac{2 \left( \frac{2 \tan x}{1 - \tan^2 x} \right)}{1 - \left( \frac{2 \tan x}{1 - \tan^2 x} \right)^2} \\ &= \frac{\left( \frac{4 \tan x}{1 - \tan^2 x} \right)}{\left[ 1 - \frac{4 \tan^2 x}{(1 - \tan^2 x)^2} \right]} \\ &= \frac{\left( \frac{4 \tan x}{1 - \tan^2 x} \right)}{\left[ \frac{(1 - \tan^2 x)^2 - 4 \tan^2 x}{(1 - \tan^2 x)^2} \right]} \\ &= \frac{4 \tan x (1 - \tan^2 x)}{(1 - \tan^2 x)^2 - 4 \tan^2 x} \\ &= \frac{4 \tan x (1 - \tan^2 x)}{1 + \tan^4 x - 2 \tan^2 x - 4 \tan^2 x} \\ &= \frac{4 \tan x (1 - \tan^2 x)}{1 - 6 \tan^2 x + \tan^4 x} = \text{R.H.S.} \end{aligned}$$

**Q24 :**

**Prove that  $\cos 4x = 1 - 8 \sin^2 x \cos^2 x$**

**Answer :**

$$\text{L.H.S.} = \cos 4x$$

$$= \cos 2(2x)$$

$$= 1 - 2 \sin^2 2x \quad [\cos 2A = 1 - 2 \sin^2 A]$$

$$= 1 - 2(2 \sin x \cos x)^2 \quad [\sin 2A = 2 \sin A \cos A]$$

$$= 1 - 8 \sin^2 x \cos^2 x$$

= R.H.S.

**Q25 :**

**Prove that:  $\cos 6x = 32 \cos^6 x - 48 \cos^4 x + 18 \cos^2 x - 1$**

**Answer :**

$$\text{L.H.S.} = \cos 6x$$

$$= \cos 3(2x)$$

$$= 4 \cos^3 2x - 3 \cos 2x [\cos 3A = 4 \cos^3 A - 3 \cos A]$$

$$= 4 [(2 \cos^2 x - 1)^3 - 3 (2 \cos^2 x - 1) [\cos 2x = 2 \cos^2 x - 1]]$$

$$= 4 [(2 \cos^2 x)^3 - (1)^3 - 3 (2 \cos^2 x)^2 + 3 (2 \cos^2 x)] - 6 \cos^2 x + 3$$

$$= 4 [8 \cos^6 x - 1 - 12 \cos^4 x + 6 \cos^2 x] - 6 \cos^2 x + 3$$

$$= 32 \cos^6 x - 4 - 48 \cos^4 x + 24 \cos^2 x - 6 \cos^2 x + 3$$

$$= 32 \cos^6 x - 48 \cos^4 x + 18 \cos^2 x - 1 =$$

R.H.S.

**Exercise 3.4 : Solutions of Questions on Page Number : 78**

**Q1 :**

**Find the principal and general solutions of the equation  $\tan x = \sqrt{3}$**

**Answer :**

$$\tan x = \sqrt{3}$$

$$\text{It is known that } \tan \frac{\pi}{3} = \sqrt{3} \text{ and } \tan \left( \frac{4\pi}{3} \right) = \tan \left( \pi + \frac{\pi}{3} \right) = \tan \frac{\pi}{3} = \sqrt{3}$$

$$\text{Therefore, the principal solutions are } x = \frac{\pi}{3} \text{ and } \frac{4\pi}{3}.$$

$$\text{Now, } \tan x = \tan \frac{\pi}{3}$$

$$\Rightarrow x = n\pi + \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}$$

$$x = n\pi + \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}$$

Therefore, the general solution is

**Q2 :**

Find the principal and general solutions of the equation  $\sec x = 2$

Answer :

$$\sec x = 2$$

It is known that  $\sec \frac{\pi}{3} = 2$  and  $\sec \frac{5\pi}{3} = \sec \left( 2\pi - \frac{\pi}{3} \right) = \sec \frac{\pi}{3} = 2$

Therefore, the principal solutions are  $x = \frac{\pi}{3}$  and  $\frac{5\pi}{3}$ .

$$\text{Now, } \sec x = \sec \frac{\pi}{3}$$

$$\Rightarrow \cos x = \cos \frac{\pi}{3} \quad \left[ \sec x = \frac{1}{\cos x} \right]$$

$$\Rightarrow x = 2n\pi \pm \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}$$

Therefore, the general solution is  $x = 2n\pi \pm \frac{\pi}{3}$ , where  $n \in \mathbb{Z}$

Q3 :

Find the principal and general solutions of the equation  $\cot x = -\sqrt{3}$

Answer :

$$\cot x = -\sqrt{3}$$

It is known that  $\cot \frac{\pi}{6} = \sqrt{3}$

$$\therefore \cot \left( \pi - \frac{\pi}{6} \right) = -\cot \frac{\pi}{6} = -\sqrt{3} \text{ and } \cot \left( 2\pi - \frac{\pi}{6} \right) = -\cot \frac{\pi}{6} = -\sqrt{3}$$

$$\text{i.e., } \cot \frac{5\pi}{6} = -\sqrt{3} \text{ and } \cot \frac{11\pi}{6} = -\sqrt{3}$$

Therefore, the principal solutions are  $x = \frac{5\pi}{6}$  and  $\frac{11\pi}{6}$ .

$$\text{Now, } \cot x = \cot \frac{5\pi}{6}$$

$$\Rightarrow \tan x = \tan \frac{5\pi}{6} \quad \left[ \cot x = \frac{1}{\tan x} \right]$$

$$\Rightarrow x = n\pi + \frac{5\pi}{6}, \text{ where } n \in \mathbb{Z}$$

$$x = n\pi + \frac{5\pi}{6}, \text{ where } n \in \mathbb{Z}$$

Therefore, the general solution is

**Q4 :**

**Find the general solution of  $\operatorname{cosec} x = -2$**

**Answer :**

$\operatorname{cosec} x = \frac{1}{\sin x}$

It is known that

$$\operatorname{cosec} \frac{\pi}{6} = 2$$

$$\therefore \operatorname{cosec} \left( \pi + \frac{\pi}{6} \right) = -\operatorname{cosec} \frac{\pi}{6} = -2 \text{ and } \operatorname{cosec} \left( 2\pi - \frac{\pi}{6} \right) = -\operatorname{cosec} \frac{\pi}{6} = -2$$

$$\text{i.e., } \operatorname{cosec} \frac{7\pi}{6} = -2 \text{ and } \operatorname{cosec} \frac{11\pi}{6} = -2$$

Therefore, the principal solutions are  $x = \frac{7\pi}{6}$  and  $\frac{11\pi}{6}$ .

$$\text{Now, } \operatorname{cosec} x = \operatorname{cosec} \frac{7\pi}{6}$$

$$\Rightarrow \sin x = \sin \frac{7\pi}{6} \quad \left[ \operatorname{cosec} x = \frac{1}{\sin x} \right]$$

$$\Rightarrow x = n\pi + (-1)^n \frac{7\pi}{6}, \text{ where } n \in \mathbb{Z}$$

$$x = n\pi + (-1)^n \frac{7\pi}{6}, \text{ where } n \in \mathbb{Z}$$

Therefore, the general solution is

**Q5 :**

**Find the general solution of the equation  $\cos 4x = \cos 2x$**

**Answer :**

$$\cos 4x = \cos 2x$$

$$\Rightarrow \cos 4x - \cos 2x = 0$$

$$\Rightarrow -2 \sin\left(\frac{4x+2x}{2}\right) \sin\left(\frac{4x-2x}{2}\right) = 0$$

$$\left[ \because \cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right) \right]$$

$$\Rightarrow \sin 3x \sin x = 0$$

$$\Rightarrow \sin 3x = 0 \quad \text{or} \quad \sin x = 0$$

$$\therefore 3x = n\pi \quad \text{or} \quad x = n\pi, \text{ where } n \in \mathbb{Z}$$

$$\Rightarrow x = \frac{n\pi}{3} \quad \text{or} \quad x = n\pi, \text{ where } n \in \mathbb{Z}$$

**Q6 :**

Find the general solution of the equation  $\cos 3x + \cos x - \cos 2x = 0$

**Answer :**

$$\cos 3x + \cos x - \cos 2x = 0$$

$$\Rightarrow 2 \cos\left(\frac{3x+x}{2}\right) \cos\left(\frac{3x-x}{2}\right) - \cos 2x = 0 \quad \left[ \cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) \right]$$

$$\Rightarrow 2 \cos 2x \cos x - \cos 2x = 0$$

$$\Rightarrow \cos 2x (2 \cos x - 1) = 0$$

$$\Rightarrow \cos 2x = 0 \quad \text{or} \quad 2 \cos x - 1 = 0$$

$$\Rightarrow \cos 2x = 0 \quad \text{or} \quad \cos x = \frac{1}{2}$$

$$\therefore 2x = (2n+1)\frac{\pi}{2} \quad \text{or} \quad \cos x = \cos \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}$$

$$\Rightarrow x = (2n+1)\frac{\pi}{4} \quad \text{or} \quad x = 2n\pi \pm \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}$$

**Q7 :**

Find the general solution of the equation  $\sin 2x + \cos x = 0$

**Answer :**

$$\sin 2x + \cos x = 0$$

$$\Rightarrow 2 \sin x \cos x + \cos x = 0$$

$$\Rightarrow \cos x (2 \sin x + 1) = 0$$

$$\Rightarrow \cos x = 0 \quad \text{or} \quad 2 \sin x + 1 = 0$$

$$\text{Now, } \cos x = 0 \Rightarrow \cos x = (2n+1) \frac{\pi}{2}, \text{ where } n \in Z$$

$$2 \sin x + 1 = 0$$

$$\Rightarrow \sin x = \frac{-1}{2} = -\sin \frac{\pi}{6} = \sin \left( \pi + \frac{\pi}{6} \right) = \sin \left( \pi + \frac{\pi}{6} \right) = \sin \frac{7\pi}{6}$$

$$\Rightarrow x = n\pi + (-1)^n \frac{7\pi}{6}, \text{ where } n \in Z$$

$$(2n+1) \frac{\pi}{2} \text{ or } n\pi + (-1)^n \frac{7\pi}{6}, n \in Z$$

Therefore, the general solution is

**Q8 :**

Find the general solution of the equation  $\sec^2 2x = 1 - \tan 2x$

**Answer :**

$$\sec^2 2x = 1 - \tan 2x$$

$$\Rightarrow 1 + \tan^2 2x = 1 - \tan 2x$$

$$\Rightarrow \tan^2 2x + \tan 2x = 0$$

$$\Rightarrow \tan 2x (\tan 2x + 1) = 0$$

$$\Rightarrow \tan 2x = 0 \quad \text{or} \quad \tan 2x + 1 = 0$$

$$\text{Now, } \tan 2x = 0$$

$$\Rightarrow \tan 2x = \tan 0$$

$$\Rightarrow 2x = n\pi + 0, \text{ where } n \in Z$$

$$\Rightarrow x = \frac{n\pi}{2}, \text{ where } n \in Z$$

$$\tan 2x + 1 = 0$$

$$\Rightarrow \tan 2x = -1 = -\tan \frac{\pi}{4} = \tan \left( \pi - \frac{\pi}{4} \right) = \tan \frac{3\pi}{4}$$

$$\Rightarrow 2x = n\pi + \frac{3\pi}{4}, \text{ where } n \in \mathbb{Z}$$

$$\Rightarrow x = \frac{n\pi}{2} + \frac{3\pi}{8}, \text{ where } n \in \mathbb{Z}$$

Therefore, the general solution is  $\frac{n\pi}{2}$  or  $\frac{n\pi}{2} + \frac{3\pi}{8}, n \in \mathbb{Z}$ .

**Q9 :**

Find the general solution of the equation  $\sin x + \sin 3x + \sin 5x = 0$

**Answer :**

$$\sin x + \sin 3x + \sin 5x = 0$$

$$(\sin x + \sin 5x) + \sin 3x = 0$$

$$\Rightarrow \left[ 2 \sin \left( \frac{x+5x}{2} \right) \cos \left( \frac{x-5x}{2} \right) \right] + \sin 3x = 0 \quad \left[ \sin A + \sin B = 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right) \right]$$

$$\Rightarrow 2 \sin 3x \cos(-2x) + \sin 3x = 0$$

$$\Rightarrow 2 \sin 3x \cos 2x + \sin 3x = 0$$

$$\Rightarrow \sin 3x (2 \cos 2x + 1) = 0$$

$$\Rightarrow \sin 3x = 0 \quad \text{or} \quad 2 \cos 2x + 1 = 0$$

Now,  $\sin 3x = 0 \Rightarrow 3x = n\pi$ , where  $n \in \mathbb{Z}$

i.e.,  $x = \frac{n\pi}{3}$ , where  $n \in \mathbb{Z}$

$$2 \cos 2x + 1 = 0$$

$$\Rightarrow \cos 2x = \frac{-1}{2} = -\cos \frac{\pi}{3} = \cos \left( \pi - \frac{\pi}{3} \right)$$

$$\Rightarrow \cos 2x = \cos \frac{2\pi}{3}$$

$$\Rightarrow 2x = 2n\pi \pm \frac{2\pi}{3}, \text{ where } n \in \mathbb{Z}$$

$$\Rightarrow x = n\pi \pm \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}$$



**Exercise Miscellaneous : Solutions of Questions on Page Number : 81****Q1 :**

Therefore, the general solution is  $\frac{n\pi}{3}$  or  $n\pi \pm \frac{\pi}{3}$ ,  $n \in \mathbb{Z}$

**Prove that:**  $2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + \cos \frac{3\pi}{13} + \cos \frac{5\pi}{13} = 0$

**Answer :**

L.H.S.

$$\begin{aligned}
 &= 2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + \cos \frac{3\pi}{13} + \cos \frac{5\pi}{13} \\
 &= 2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + 2 \cos \left( \frac{\frac{3\pi}{13} + \frac{5\pi}{13}}{2} \right) \cos \left( \frac{\frac{3\pi}{13} - \frac{5\pi}{13}}{2} \right) \left[ \cos x + \cos y = 2 \cos \left( \frac{x+y}{2} \right) \cos \left( \frac{x-y}{2} \right) \right] \\
 &= 2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + 2 \cos \frac{4\pi}{13} \cos \left( \frac{-\pi}{13} \right) \\
 &= 2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + 2 \cos \frac{4\pi}{13} \cos \frac{\pi}{13} \\
 &= 2 \cos \frac{\pi}{13} \left[ \cos \frac{9\pi}{13} + \cos \frac{4\pi}{13} \right] \\
 &= 2 \cos \frac{\pi}{13} \left[ 2 \cos \left( \frac{\frac{9\pi}{13} + \frac{4\pi}{13}}{2} \right) \cos \left( \frac{\frac{9\pi}{13} - \frac{4\pi}{13}}{2} \right) \right] \\
 &= 2 \cos \frac{\pi}{13} \left[ 2 \cos \frac{\pi}{2} \cos \frac{5\pi}{26} \right] \\
 &= 2 \cos \frac{\pi}{13} \times 2 \times 0 \times \cos \frac{5\pi}{26}
 \end{aligned}$$

 $= 0 = \text{R.H.S}$ **Q2 :****Prove that:**  $(\sin 3x + \sin x) \sin x + (\cos 3x - \cos x) \cos x = 0$ **Answer :**

L.H.S.

$$= (\sin 3x + \sin x) \sin x + (\cos 3x - \cos x) \cos x$$

$$= \sin 3x \sin x + \sin^2 x + \cos 3x \cos x - \cos^2 x$$

$$= \cos 3x \cos x + \sin 3x \sin x - (\cos^2 x - \sin^2 x)$$

$$= \cos(3x - x) - \cos 2x \quad [\cos(A - B) = \cos A \cos B + \sin A \sin B]$$

$$= \cos 2x - \cos 2x$$

$$= 0$$

= R.H.S.

**Q3 :**

Prove that:  $(\cos x + \cos y)^2 + (\sin x - \sin y)^2 = 4 \cos^2 \frac{x+y}{2}$

**Answer :**

$$\text{L.H.S.} = (\cos x + \cos y)^2 + (\sin x - \sin y)^2$$

$$= \cos^2 x + \cos^2 y + 2 \cos x \cos y + \sin^2 x + \sin^2 y - 2 \sin x \sin y$$

$$= (\cos^2 x + \sin^2 x) + (\cos^2 y + \sin^2 y) + 2(\cos x \cos y - \sin x \sin y)$$

$$= 1 + 1 + 2 \cos(x + y) \quad [\cos(A + B) = (\cos A \cos B - \sin A \sin B)]$$

$$= 2 + 2 \cos(x + y)$$

$$= 2[1 + \cos(x + y)]$$

$$= 2 \left[ 1 + 2 \cos^2 \left( \frac{x+y}{2} \right) - 1 \right] \quad [\cos 2A = 2 \cos^2 A - 1]$$

$$= 4 \cos^2 \left( \frac{x+y}{2} \right) = \text{R.H.S.}$$

**Q4 :**

Prove that:  $(\cos x - \cos y)^2 + (\sin x - \sin y)^2 = 4 \sin^2 \frac{x-y}{2}$

**Answer :**

$$\text{L.H.S.} = (\cos x - \cos y)^2 + (\sin x - \sin y)^2$$

$$\begin{aligned}
&= \cos^2 x + \cos^2 y - 2 \cos x \cos y + \sin^2 x + \sin^2 y - 2 \sin x \sin y \\
&= (\cos^2 x + \sin^2 x) + (\cos^2 y + \sin^2 y) - 2[\cos x \cos y + \sin x \sin y] \\
&= 1 + 1 - 2[\cos(x - y)] \quad [\cos(A - B) = \cos A \cos B + \sin A \sin B] \\
&= 2[1 - \cos(x - y)] \\
&= 2 \left[ 1 - \left\{ 1 - 2 \sin^2 \left( \frac{x - y}{2} \right) \right\} \right] \quad [\cos 2A = 1 - 2 \sin^2 A] \\
&= 4 \sin^2 \left( \frac{x - y}{2} \right) = \text{R.H.S.}
\end{aligned}$$

**Q5 :**

**Prove that:**  $\sin x + \sin 3x + \sin 5x + \sin 7x = 4 \cos x \cos 2x \sin 4x$

**Answer :**

$$\sin A + \sin B = 2 \sin \left( \frac{A + B}{2} \right) \cdot \cos \left( \frac{A - B}{2} \right)$$

It is known that

$$\therefore \text{L.H.S.} = \sin x + \sin 3x + \sin 5x + \sin 7x$$

$$\begin{aligned}
&= (\sin x + \sin 5x) + (\sin 3x + \sin 7x) \\
&= 2 \sin \left( \frac{x + 5x}{2} \right) \cdot \cos \left( \frac{x - 5x}{2} \right) + 2 \sin \left( \frac{3x + 7x}{2} \right) \cos \left( \frac{3x - 7x}{2} \right) \\
&= 2 \sin 3x \cos(-2x) + 2 \sin 5x \cos(-2x) \\
&= 2 \sin 3x \cos 2x + 2 \sin 5x \cos 2x \\
&= 2 \cos 2x [\sin 3x + \sin 5x] \\
&= 2 \cos 2x \left[ 2 \sin \left( \frac{3x + 5x}{2} \right) \cdot \cos \left( \frac{3x - 5x}{2} \right) \right] \\
&= 2 \cos 2x [2 \sin 4x \cdot \cos(-x)] \\
&= 4 \cos 2x \sin 4x \cos x = \text{R.H.S.}
\end{aligned}$$

**Q6 :**

$$\text{Prove that: } \frac{(\sin 7x + \sin 5x) + (\sin 9x + \sin 3x)}{(\cos 7x + \cos 5x) + (\cos 9x + \cos 3x)} = \tan 6x$$

**Answer :**

It is known that

$$\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cdot \cos\left(\frac{A-B}{2}\right), \quad \cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cdot \cos\left(\frac{A-B}{2}\right)$$

$$\text{L.H.S.} = \frac{(\sin 7x + \sin 5x) + (\sin 9x + \sin 3x)}{(\cos 7x + \cos 5x) + (\cos 9x + \cos 3x)}$$

$$\begin{aligned} &= \frac{\left[2 \sin\left(\frac{7x+5x}{2}\right) \cdot \cos\left(\frac{7x-5x}{2}\right)\right] + \left[2 \sin\left(\frac{9x+3x}{2}\right) \cdot \cos\left(\frac{9x-3x}{2}\right)\right]}{\left[2 \cos\left(\frac{7x+5x}{2}\right) \cdot \cos\left(\frac{7x-5x}{2}\right)\right] + \left[2 \cos\left(\frac{9x+3x}{2}\right) \cdot \cos\left(\frac{9x-3x}{2}\right)\right]} \\ &= \frac{[2 \sin 6x \cdot \cos x] + [2 \sin 6x \cdot \cos 3x]}{[2 \cos 6x \cdot \cos x] + [2 \cos 6x \cdot \cos 3x]} \\ &= \frac{2 \sin 6x [\cos x + \cos 3x]}{2 \cos 6x [\cos x + \cos 3x]} \end{aligned}$$

$$= \tan 6x$$

$$= \text{R.H.S.}$$

**Q7 :**

$$\sin 3x + \sin 2x - \sin x = 4 \sin x \cos \frac{x}{2} \cos \frac{3x}{2}$$

**Prove that:**

**Answer :**

$$\text{L.H.S.} = \sin 3x + \sin 2x - \sin x$$

$$\begin{aligned}
&= \sin 3x + (\sin 2x - \sin x) \\
&= \sin 3x + \left[ 2 \cos \left( \frac{2x+x}{2} \right) \sin \left( \frac{2x-x}{2} \right) \right] \quad \left[ \sin A - \sin B = 2 \cos \left( \frac{A+B}{2} \right) \sin \left( \frac{A-B}{2} \right) \right] \\
&= \sin 3x + \left[ 2 \cos \left( \frac{3x}{2} \right) \sin \left( \frac{x}{2} \right) \right] \\
&= \sin 3x + 2 \cos \frac{3x}{2} \sin \frac{x}{2} \\
&= 2 \sin \frac{3x}{2} \cdot \cos \frac{3x}{2} + 2 \cos \frac{3x}{2} \sin \frac{x}{2} \quad [\sin 2A = 2 \sin A \cdot \cos B] \\
&= 2 \cos \left( \frac{3x}{2} \right) \left[ \sin \left( \frac{3x}{2} \right) + \sin \left( \frac{x}{2} \right) \right] \\
&= 2 \cos \left( \frac{3x}{2} \right) \left[ 2 \sin \left\{ \frac{\left( \frac{3x}{2} \right) + \left( \frac{x}{2} \right)}{2} \right\} \cos \left\{ \frac{\left( \frac{3x}{2} \right) - \left( \frac{x}{2} \right)}{2} \right\} \right] \quad \left[ \sin A + \sin B = 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right) \right] \\
&= 2 \cos \left( \frac{3x}{2} \right) \cdot 2 \sin x \cos \left( \frac{x}{2} \right) \\
&= 4 \sin x \cos \left( \frac{x}{2} \right) \cos \left( \frac{3x}{2} \right) = \text{R.H.S.}
\end{aligned}$$

**Q8 :**

$$\tan x = -\frac{4}{3}, \text{ } x \text{ in quadrant II}$$

**Answer :**

Here,  $x$  is in quadrant II.

$$\text{i.e., } \frac{\pi}{2} < x < \pi$$

$$\Rightarrow \frac{\pi}{4} < \frac{x}{2} < \frac{\pi}{2}$$

Therefore,  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$  and  $\tan \frac{x}{2}$  are all positive.

It is given that  $\tan x = -\frac{4}{3}$ .

$$\sec^2 x = 1 + \tan^2 x = 1 + \left(\frac{-4}{3}\right)^2 = 1 + \frac{16}{9} = \frac{25}{9}$$

$$\therefore \cos^2 x = \frac{9}{25}$$

$$\Rightarrow \cos x = \pm \frac{3}{5}$$

As  $x$  is in quadrant II,  $\cos x$  is negative.

$$\therefore \cos x = -\frac{3}{5}$$

$$\text{Now, } \cos x = 2 \cos^2 \frac{x}{2} - 1$$

$$\Rightarrow \frac{-3}{5} = 2 \cos^2 \frac{x}{2} - 1$$

$$\Rightarrow 2 \cos^2 \frac{x}{2} = 1 - \frac{3}{5}$$

$$\Rightarrow 2 \cos^2 \frac{x}{2} = \frac{2}{5}$$

$$\Rightarrow \cos^2 \frac{x}{2} = \frac{1}{5}$$

$$\Rightarrow \cos \frac{x}{2} = \frac{1}{\sqrt{5}} \quad \left[ \because \cos \frac{x}{2} \text{ is positive} \right]$$

$$\therefore \cos \frac{x}{2} = \frac{\sqrt{5}}{5}$$

$$\sin^2 \frac{x}{2} + \cos^2 \frac{x}{2} = 1$$

$$\Rightarrow \sin^2 \frac{x}{2} + \left(\frac{1}{\sqrt{5}}\right)^2 = 1$$

$$\Rightarrow \sin^2 \frac{x}{2} = 1 - \frac{1}{5} = \frac{4}{5}$$

$$\Rightarrow \sin \frac{x}{2} = \frac{2}{\sqrt{5}} \quad \left[ \because \sin \frac{x}{2} \text{ is positive} \right]$$

$$\therefore \sin \frac{x}{2} = \frac{2\sqrt{5}}{5}$$

$$\tan \frac{x}{2} = \frac{\sin \frac{x}{2}}{\cos \frac{x}{2}} = \frac{\left(\frac{2}{\sqrt{5}}\right)}{\left(\frac{1}{\sqrt{5}}\right)} = 2$$

Thus, the respective values of  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$  and  $\tan \frac{x}{2}$  are  $\frac{2\sqrt{5}}{5}$ ,  $\frac{\sqrt{5}}{5}$ , and 2.

**Q9 :**

Find  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$  and  $\tan \frac{x}{2}$  for  $\cos x = -\frac{1}{3}$ ,  $x$  in quadrant III

**Answer :**

Here,  $x$  is in quadrant III.

$$\text{i.e., } \pi < x < \frac{3\pi}{2}$$

$$\Rightarrow \frac{\pi}{2} < \frac{x}{2} < \frac{3\pi}{4}$$

Therefore,  $\cos \frac{x}{2}$  and  $\tan \frac{x}{2}$  are negative, whereas  $\sin \frac{x}{2}$  is positive.

It is given that  $\cos x = -\frac{1}{3}$ .

$$\cos x = 1 - 2 \sin^2 \frac{x}{2}$$

$$\Rightarrow \sin^2 \frac{x}{2} = \frac{1 - \cos x}{2}$$

$$\Rightarrow \sin^2 \frac{x}{2} = \frac{1 - \left(-\frac{1}{3}\right)}{2} = \frac{\left(1 + \frac{1}{3}\right)}{2} = \frac{\frac{4}{3}}{2} = \frac{2}{3}$$

$$\Rightarrow \sin \frac{x}{2} = \frac{\sqrt{2}}{\sqrt{3}} \quad \left[ \because \sin \frac{x}{2} \text{ is positive} \right]$$

$$\therefore \sin \frac{x}{2} = \frac{\sqrt{2}}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{\sqrt{6}}{3}$$

$$\cos x = 2 \cos^2 \frac{x}{2} - 1$$

Now,

$$\Rightarrow \cos^2 \frac{x}{2} = \frac{1 + \cos x}{2} = \frac{1 + \left(-\frac{1}{3}\right)}{2} = \frac{\left(\frac{3-1}{3}\right)}{2} = \frac{\left(\frac{2}{3}\right)}{2} = \frac{1}{3}$$

$$\Rightarrow \cos \frac{x}{2} = -\frac{1}{\sqrt{3}} \quad \left[ \because \cos \frac{x}{2} \text{ is negative} \right]$$

$$\therefore \cos \frac{x}{2} = -\frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{-\sqrt{3}}{3}$$

$$\tan \frac{x}{2} = \frac{\sin \frac{x}{2}}{\cos \frac{x}{2}} = \frac{\left(\frac{\sqrt{2}}{\sqrt{3}}\right)}{\left(\frac{-1}{\sqrt{3}}\right)} = -\sqrt{2}$$

Thus, the respective values of  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$  and  $\tan \frac{x}{2}$  are  $\frac{\sqrt{6}}{3}$ ,  $\frac{-\sqrt{3}}{3}$ , and  $-\sqrt{2}$ .

**Q10 :**

Find  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$  and  $\tan \frac{x}{2}$  for  $\sin x = \frac{1}{4}$ ,  $x$  in quadrant II

**Answer :**

Here,  $x$  is in quadrant II.

$$\text{i.e., } \frac{\pi}{2} < x < \pi$$

$$\Rightarrow \frac{\pi}{4} < \frac{x}{2} < \frac{\pi}{2}$$

Therefore,  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$ , and  $\tan \frac{x}{2}$  are all positive.

It is given that  $\sin x = \frac{1}{4}$ .

$$\cos^2 x = 1 - \sin^2 x = 1 - \left(\frac{1}{4}\right)^2 = 1 - \frac{1}{16} = \frac{15}{16}$$

$$\Rightarrow \cos x = -\frac{\sqrt{15}}{4} \quad [\cos x \text{ is negative in quadrant II}]$$



$$\sin^2 \frac{x}{2} = \frac{1 - \cos x}{2} = \frac{1 - \left(-\frac{\sqrt{15}}{4}\right)}{2} = \frac{4 + \sqrt{15}}{8}$$

$$\Rightarrow \sin \frac{x}{2} = \sqrt{\frac{4 + \sqrt{15}}{8}} \quad \left[ \because \sin \frac{x}{2} \text{ is positive} \right]$$

$$= \sqrt{\frac{4 + \sqrt{15}}{8}} \times \frac{2}{2}$$

$$= \sqrt{\frac{8 + 2\sqrt{15}}{16}}$$

$$= \frac{\sqrt{8 + 2\sqrt{15}}}{4}$$

$$\cos^2 \frac{x}{2} = \frac{1 + \cos x}{2} = \frac{1 + \left(-\frac{\sqrt{15}}{4}\right)}{2} = \frac{4 - \sqrt{15}}{8}$$

$$\Rightarrow \cos \frac{x}{2} = \sqrt{\frac{4 - \sqrt{15}}{8}} \quad \left[ \because \cos \frac{x}{2} \text{ is positive} \right]$$

$$= \sqrt{\frac{4 - \sqrt{15}}{8}} \times \frac{2}{2}$$

$$= \sqrt{\frac{8 - 2\sqrt{15}}{16}}$$

$$= \frac{\sqrt{8 - 2\sqrt{15}}}{4}$$

$$\tan \frac{x}{2} = \frac{\sin \frac{x}{2}}{\cos \frac{x}{2}} = \frac{\left(\frac{\sqrt{8 + 2\sqrt{15}}}{4}\right)}{\left(\frac{\sqrt{8 - 2\sqrt{15}}}{4}\right)} = \frac{\sqrt{8 + 2\sqrt{15}}}{\sqrt{8 - 2\sqrt{15}}}$$

$$= \sqrt{\frac{8 + 2\sqrt{15}}{8 - 2\sqrt{15}}} \times \frac{8 + 2\sqrt{15}}{8 + 2\sqrt{15}}$$

$$= \sqrt{\frac{(8 + 2\sqrt{15})^2}{64 - 60}} = \frac{8 + 2\sqrt{15}}{2} = 4 + \sqrt{15}$$

Thus, the respective values of  $\sin \frac{x}{2}$ ,  $\cos \frac{x}{2}$  and  $\tan \frac{x}{2}$  are  $\frac{\sqrt{8+2\sqrt{15}}}{4}$ ,  $\frac{\sqrt{8-2\sqrt{15}}}{4}$ ,  
and  $4 + \sqrt{15}$ .